

On the Canonical Reduction of Spherically Symmetric Gravity*

Stephen R. Lau[†]

*Institut für Theoretische Physik
Technische Universität Wien
Wiedner Hauptstraße 8-10
A-1040 Wien, AUSTRIA*

Abstract

In a thorough paper Kuchař has examined the canonical reduction of the most general action functional describing the geometrodynamics of the maximally extended Schwarzschild geometry. This reduction yields the true degrees of freedom for (vacuum) spherically symmetric general relativity (SSGR). The essential technical ingredient in Kuchař’s analysis is a canonical transformation to a certain chart on the gravitational phase space which features the Schwarzschild mass parameter M_S , expressed in terms of what are essentially Arnowitt-Deser-Misner variables, as a canonical coordinate. (Kuchař’s paper complements earlier work by Kastrup and Thiemann, based mostly on Ashtekar variables, which has also explicitly isolated the true degrees of freedom for vacuum SSGR.) In this paper we discuss the geometric interpretation of Kuchař’s canonical transformation in terms of the theory of quasilocal energy-momentum in general relativity given by Brown and York. We find Kuchař’s transformation to be a “sphere-dependent boost to the rest frame,” where the “rest frame” is defined by vanishing quasilocal momentum. Furthermore, our formalism is general enough to cover the case of (vacuum) two-dimensional dilaton gravity. Therefore, besides reviewing Kuchař’s original work for Schwarzschild black holes from the framework of hyperbolic geometry, we present new results concerning the canonical reduction of Witten-black-hole geometrodynamics. Finally, addressing a recent work of Louko and Whiting, we discuss some delicate points concerning the canonical reduction of the “thermodynamical action,” which is of central importance in the path-integral formulation of gravitational thermodynamics.

Vienna, February 1996

*Revised version (to appear in *Classical and Quantum Gravity*).

[†]email address: lau@thp16.tuwien.ac.at

I. INTRODUCTION

In a thorough paper (Ref. [1], hereafter referred to as KVK) Kuchař has examined the canonical reduction of the most general action functional describing the geometrodynamics of the maximally extended Schwarzschild geometry. This reduction yields the true degrees of freedom for vacuum general relativity subject to ansatz of spherical symmetry. The key technical ingredient in Kuchař’s analysis is a canonical transformation to a certain chart on the gravitational phase space which features the Schwarzschild mass parameter M_s , expressed in terms of what are essentially Arnowitt-Deser-Misner (ADM) variables, [2] as a canonical coordinate. (Kuchař’s paper complements earlier work [3] by Kastrup and Thiemann, based mostly on Ashtekar variables, which has also explicitly isolated the true degrees of freedom for vacuum SSGR.) Potential applications of the new reduced formalism include examinations, from the canonical viewpoint, of spherically symmetric collapse, the Hawking effect, and equilibrium gravitational thermodynamics. Indeed, Louko and Whiting have already made such an application. Applying Kuchař’s method to a spatially bounded exterior region of the Schwarzschild black hole, they have constructed the Schwarzschild thermodynamical (canonical) partition function completely within the Lorentzian Hamiltonian framework (Ref. [4], hereafter referred to as LW). Their canonical partition function is in agreement with previous results derived via the Euclidean-path-integral method. [5] The starting point in LW is the “thermodynamical action,” which is of central importance in the path-integral formulation of gravitational thermodynamics. A very delicate issue in the analysis of LW concerns the treatment of the thermodynamical action’s boundary terms under the canonical reduction via the KVK method.

In this paper we discuss the geometric interpretation of Kuchař’s canonical transformation. By appealing to notions of quasilocal energy and momentum in general relativity which have been given by Brown and York [6,7], we interpret Kuchař’s canonical transformation as a “sphere-dependent boost to the rest frame,” where the “rest frame” is defined by vanishing quasilocal momentum. It seems to us that KVK finds essentially the same interpretation via an alternate route involving subsequent canonical transformations and a “parameterization at infinity.” We believe that our approach complements the one given by KVK. The center-stone of ours is the following observation. On an arbitrary (spherically symmetric) spatial slice Σ the parameter φ describing the local boost between the slice Eulerian observers at a point and the rest-frame observers at the same point can be constructed from the canonical variables of Σ (if known in a tiny spatial region surrounding the point of interest). Furthermore, we work in a framework which is general enough to cover Witten’s two-dimensional dilaton gravity (2dDG). [8] (Further still, many of our preliminary results concerning the theory of quasilocal energy-momentum are unaffected by the inclusion of minimally coupled matter and hence are relevant for the two-dimensional dilaton-plus-matter model of Callan, Giddings, Harvey, and Strominger. [9]) Therefore, besides reviewing some of Kuchař’s original work for Schwarzschild black holes from the framework of hyperbolic geometry, we present new results concerning the canonical reduction of Witten-black-hole geometrodynamics. We show that the canonical transformation of KVK can also be made in the context of (vacuum) 2dDG. Therefore, the potential applications of the KVK formalism, listed in the first paragraph, are also relevant for 2dDG. Finally, with our general framework we address some of the delicate points, first considered in LW, concerning the canonical re-

duction of the “thermodynamical action.” Our conceptual framework supports the difficult technical steps taken in LW. All of our results are given for both the Schwarzschild and Witten-black-hole cases.

A few technical points demand some comment at the outset. As mentioned, the analysis of KVK concerns the full Kruskal spacetime, the maximally extended Schwarzschild geometry. The canonical variables used in KVK are defined on spatial slices which cut completely across the Kruskal diagram, and therefore have to obey appropriate boundary conditions in the asymptotic regions. In crossing from one spatial infinity to the other, the slices of KVK are allowed to cross the horizon in a completely general way. This introduces some technical difficulties at the horizon, especially when one is considering Kuchař’s canonical transformation. However, as demonstrated in KVK, with care these difficulties may be surmounted. We choose to confine our attention entirely to the right static region of the Kruskal diagram. At first, we work with the time history of a static region lying between concentric spheres. Thus we avoid many of the technical difficulties faced by Kuchař at the outset. We could, of course, work in the full Kruskal diagram, but the essential points of this paper do not demand that we do so. However, since we do chose to bypass a technical treatment of the horizon, questions concerning how to handle such horizon difficulties remain for the Witten black holes of 2dDG. However, notationally we adopt nearly the same conventions as KVK. Therefore, we expect that with the present paper as a stepping stone, the interested reader could -with minimal effort- convert any and all of the horizon arguments given in KVK into corresponding arguments applicable to the Witten black-hole case.

The layout of this paper is as follows. In § II, the preliminary section, we describe the relevant kinematics of our spacetime geometry. Since the spacetime geometry is spherically symmetric, it proves convenient to work with a toy 1+1 dimensional spacetime $\mathcal{M}$. In reality, the points of $\mathcal{M}$ are round spheres. In § III we derive quasilocal¹ energy and momentum expressions for the physical fields defined on a generic spatial slice Σ of $\mathcal{M}$. The method used to derive the quasilocal expressions is a Hamilton-Jacobi analysis of an appropriate action principle for $\mathcal{M}$. In § IV we use the developed notions of quasilocal energy and momentum to underscore the geometric significance of Kuchař’s canonical transformation. This section also considers the reduction of the canonical action with the boundary conditions adopted in this work. Finally, § V considers the canonical reduction of the thermodynamical action.

II. PRELIMINARIES

A. Spacetime $\mathcal{M}$ and foliations

Consider a 1+1 dimensional spacetime region $\mathcal{M}$ which is bounded spatially. The region $\mathcal{M}$ consists of a collection of one dimensional spacelike slices Σ . The letter Σ denotes both a foliation of $\mathcal{M}$ into spacelike slices and a generic leaf of such a foliation. However, for

¹We use the adjective “quasilocal” because from the four-dimensional viewpoint the energy and momentum are associated round two-spheres. Thus these “quasilocal” quantities are associated with points of $\mathcal{M}$.

the initial spacelike slice we reserve the special symbol t' (also the value of the coordinate time on the initial slice), and, likewise, for the final spacelike slice we reserve the symbol t'' (also the value of the coordinate time on the final slice). On spacetime $\mathcal{M}$ we have coordinates (t, r) , and a generic spacetime point² is $B(t, r)$. Every constant- t slice Σ has two boundary points B_i (at $r = r_i$) and B_o (at $r = r_o$). Assume that along Σ the coordinate r increases monotonically from B_i to B_o . We represent the timelike history $B_i(t) \equiv B(t, r_i)$ by $\bar{\mathcal{T}}_i$ (unbarred $\mathcal{T}$ is reserved for another meaning) and refer to it as the *inner boundary*. Likewise, we represent the timelike history $B_o(t) \equiv B(t, r_o)$ by $\bar{\mathcal{T}}_o$ and refer to it as the *outer boundary*. Later on, when we deal with black-hole solutions, we will “seal” the inner boundary. In other words, the time development at the inner boundary will be arrested, and the point B_i will correspond to a bifurcation point in a Kruskal-like diagram. We denote the *corner* points of our spacetime as follows: $B'_i \equiv B(t', r_i)$, $B'_o \equiv B(t', r_o)$, $B''_i \equiv B(t'', r_i)$, and $B''_o \equiv B(t'', r_o)$.

We can also consider a *radial* foliation of $\mathcal{M}$ by a family of one-dimensional timelike surfaces which extend from $\bar{\mathcal{T}}_i$ outward to $\bar{\mathcal{T}}_o$ (for black-hole solutions $\bar{\mathcal{T}}_i$ may be a degenerate sheet). [10] These are constant- r surfaces. Like before, we loosely use the letter $\bar{\mathcal{T}}$ both to denote the radial foliation and a generic leaf of this foliation. We call $\bar{\mathcal{T}}$ leaves *sheets*, whereas we have called Σ leaves *slices* (this is an informal convention). Abusing the notation a bit, we also often let the symbol $\bar{\mathcal{T}}$ denote the total timelike boundary $\bar{\mathcal{T}}_i \cup \bar{\mathcal{T}}_o$. However, when the symbol $\bar{\mathcal{T}}$ has this meaning, it always appears in the phrase “the $\bar{\mathcal{T}}$ boundary.”

B. Metric decompositions

The spacetime metric is g_{ab} . The metric on a generic Σ slice is Λ^2 , and the metric on the $\bar{\mathcal{T}}$ boundary is denoted by $-\bar{N}^2$. In terms of the Σ foliation, the metric may be written in ADM form [2]

$$g_{ab}dx^a dx^b = -N^2 dt^2 + \Lambda^2 (dr + N^r dt)^2, \quad (2.1)$$

with N and N^r denoting the familiar *lapse* and *shift*. The vector field

$$u^a \partial / \partial x^a = N^{-1} (\partial / \partial t - N^r \partial / \partial r) \quad (2.2)$$

is the unit, timelike, future-pointing normal to the Σ foliation.

In terms of the $\bar{\mathcal{T}}$ foliation, the $\mathcal{M}$ metric takes the form

$$g_{ab}dx^a dx^b = \bar{\Lambda}^2 dr^2 - \bar{N}^2 (dt + \bar{\Lambda}^t dr)^2, \quad (2.3)$$

where $\bar{\Lambda}$ and $\bar{\Lambda}^t$ are the *radial lapse* and *radial shift*. The unit, spacelike, $\bar{\mathcal{T}}$ -foliation normal is

²One may consider each point of $\mathcal{M}$ to be round two-sphere with radius $R(t, r)$, where R is the radius function.

$$\bar{n}^a \partial / \partial x^a = \bar{\Lambda}^{-1} \left(\partial / \partial r - \bar{\Lambda}^t \partial / \partial t \right). \quad (2.4)$$

We also define the $\bar{\mathcal{T}}$ boundary normal $\bar{n}^a$ on $\bar{\mathcal{T}}_i$ and $\bar{\mathcal{T}}_o$ by the requirement that it always be *outward*-pointing on these boundary elements. On the outer boundary $\bar{\mathcal{T}}_o$ the outward normal is $\bar{n}^a = \bar{\mathbf{n}}^a$, while on the inner boundary $\bar{\mathcal{T}}_i$ the outward normal is $\bar{n}^a = -\bar{\mathbf{n}}^a$. (For a handful $\{n^a, \bar{n}^a, v, w, \eta, \varphi, E\}$ of symbols we use this convention throughout the paper. Regular letters represent objects associated with the boundary and have the appropriate sign for each boundary element built in. Boldface versions of the same letters represent the same objects but with the fixed sign appropriate for the outer boundary. Perhaps it would have been a more natural choice to adopt the reciprocal notation and let the boldface letters possess the sign flexibility. We have made the seemingly unnatural choice simply because it leads to the minimal use of boldface letters. In other words, in what follows the sign-flexible quantities happen to appear more frequently than their fixed-sign counterparts.)

By equating the coefficients of the above forms of g_{ab} , we obtain the following relations between the “barred” and “unbarred” variables:

$$\bar{N} = N/\gamma \quad (2.5a)$$

$$\bar{\Lambda} = \gamma \Lambda \quad (2.5b)$$

$$\Lambda N^r / N = -\bar{N} \bar{\Lambda}^t / \bar{\Lambda}. \quad (2.5c)$$

Here $\gamma \equiv (1 - \mathbf{v}^2)^{-1/2}$ is the local relativistic factor associated with the velocity $\mathbf{v} \equiv \Lambda N^r / N = -\bar{N} \bar{\Lambda}^t / \bar{\Lambda} = -\bar{v}$. [10] The timelike normal associated with the foliations $B_i(t)$ and $B_o(t)$ of the boundary sheets $\bar{\mathcal{T}}_i$ and $\bar{\mathcal{T}}_o$ is $\bar{u}^a \partial / \partial x^a = \bar{N}^{-1} \partial / \partial t$. Note that on the $\bar{\mathcal{T}}$ boundary the vector fields u^a and $\bar{u}^a$ need not coincide. Also, fixation of the t coordinate gives a collection of points $B(r)$ which foliates the slice Σ . The normal associated with this foliation of Σ is $\mathbf{n}^a \partial / \partial x^a = \Lambda^{-1} \partial / \partial r$. Again, we define a boundary normal n^a such that at the inner boundary n^a is $-\mathbf{n}^a$, the outward-pointing normal of B_i as embedded in Σ , while at the outer boundary n^a is $\mathbf{n}^a$, the outward-pointing normal of B_o as embedded in Σ . On the inner and outer boundaries n^a and $\bar{n}^a$ need not coincide. It is easy to verify the following point-wise boost relations:

$$\bar{u}^a = \gamma u^a + v \gamma n^a \quad (2.6a)$$

$$\bar{n}^a = \gamma n^a + v \gamma u^a, \quad (2.6b)$$

where $v = -\mathbf{v}$ on $\bar{\mathcal{T}}_i$ and $v = \mathbf{v}$ on $\bar{\mathcal{T}}_o$.

C. Extrinsic curvatures

The extrinsic curvature of a Σ slice as embedded in $\mathcal{M}$ is $\mathcal{K} \equiv -\nabla_a u^a$, where ∇_a denotes the torsion-free covariant derivative operator compatible with g_{ab} . One may also consider spacelike slices $\bar{\Sigma}$ which are everywhere orthogonal by assumption to the $\bar{\mathcal{T}}$ sheets. Since the spacetime-filling extension of the $\bar{\mathcal{T}}$ sheets from $\bar{\mathcal{T}}_i$ to $\bar{\mathcal{T}}_o$ is arbitrary, the $\bar{\Sigma}$ slices are almost as general as the Σ slices. However, the $\bar{\Sigma}$ slices are restricted by the requirement that their normal vector field coincides with $\bar{u}^a$ on the $\bar{\mathcal{T}}$ boundary. We describe such slices as *clamped*. When the velocity v defined above is set to zero on the boundary, then the Σ

slices are clamped. (In which case, there is no longer a need to make a distinction between barred and unbarred slices.) Also define $\bar{\mathcal{K}} \equiv -\nabla_a \bar{u}^a$, the extrinsic curvature associated with the $\bar{\Sigma}$ slices. The extrinsic curvature associated with the $\bar{\mathcal{T}}$ boundary elements is defined by $\bar{\vartheta} \equiv -\nabla_a \bar{n}^a$. We may also consider a foliation $\mathcal{T}$ generated by the u^a Eulerian histories of points in the Σ slices. At the boundary, the $\mathcal{T}$ sheets may be “crashing into” or “emerging from” the actual boundary elements $\bar{\mathcal{T}}_i$ and $\bar{\mathcal{T}}_o$. Nevertheless, one can define an extrinsic curvature $\vartheta \equiv -\nabla_a n^a$. The value of ϑ at a particular boundary point of the $\bar{\mathcal{T}}$ boundary is associated with the $\mathcal{T}$ sheet intersecting that point. We have that $\vartheta = -n_b a^b$, where $a^b \equiv u^a \nabla_a u^b$ is the spacetime acceleration of u^a . Since by assumption the metric-compatible connection associated with $\mathcal{M}$ is torsion-free, $-\vartheta = n^a \nabla_a [\log N]$.

With our transformation equations (2.6), one can derive the following “splitting” formulae for $\bar{\mathcal{K}}$ and $\bar{\vartheta}$:

$$\bar{\mathcal{K}} = \gamma \mathcal{K} + v \gamma \vartheta - \bar{n}^a \nabla_a \eta \quad (2.7a)$$

$$\bar{\vartheta} = \gamma \vartheta + v \gamma \mathcal{K} - \bar{u}^a \nabla_a \eta, \quad (2.7b)$$

where $\eta \equiv \tanh^{-1} v = \frac{1}{2} \log |(1+v)/(1-v)|$. Note that $\eta = \tanh^{-1} -\boldsymbol{v}$ on $\bar{\mathcal{T}}_i$ and $\eta = \tanh^{-1} \boldsymbol{v}$ on $\bar{\mathcal{T}}_o$. In accord with our conventions also set $\boldsymbol{\eta} \equiv \tanh^{-1} \boldsymbol{v}$.

III. ACTION AND QUASILOCAL ENERGY-MOMENTUM

We begin this section by precisely defining the type of action functional associated with our bounded spacetime region $\mathcal{M}$ which is of interest in this work. We will discuss in detail the action’s associated variational principle, paying strict attention to all boundary terms. This is the background work necessary to derive expressions for the quasilocal energy and momentum associated with a generic bounded slice Σ .

A. Variational principle

Our analysis begins with the following action functional:

$$\begin{aligned} \mathcal{S}^I = & \frac{1}{4} \alpha \int_{\mathcal{M}} d^2x \sqrt{-g} e^{-2\Phi} \left[\mathcal{R} + 2y g^{ab} \nabla_a \Phi \nabla_b \Phi + 2\lambda^2 y \exp[(2-y)2\Phi] \right] \\ & + \frac{1}{2} \alpha \int_{t''}^{t'} dr \Lambda e^{-2\Phi} \mathcal{K} - \frac{1}{2} \alpha \int_{\bar{\mathcal{T}}} dt \bar{N} e^{-2\Phi} \bar{\vartheta} - \frac{1}{2} \alpha e^{-2\Phi} \eta \Big|_{B'}^{B''}, \end{aligned} \quad (3.1)$$

where $\mathcal{R}$ is the scalar curvature of $\mathcal{M}$ built from the metric g_{ab} , and the scalar field Φ is the celebrated *dilaton*. The variable y is an as-yet unspecified number, λ is a positive constant with dimensions of inverse length, and α is another positive (and possibly dimensionful) constant. The $\mathcal{M}$ integral in our action corresponds to a subclass of models within the larger framework of generalized dilaton theories. [11] In (3.1) and throughout the rest of this paper we use the following short-hand notations:

$$\int_{t'}^{t''} = \int_{t''} - \int_{t'} \quad (3.2a)$$

$$\int_{\bar{T}} = \int_{\bar{T}_i} + \int_{\bar{T}_o} \quad (3.2b)$$

$$\left|_{B'}^{B''} = \left|_{B''_i} - \left|_{B'_i} + \left|_{B''_o} - \left|_{B'_o} \right. \right. , \quad (3.2c)$$

where (3.2a) is used only when t' and t'' represent the initial and final slices and not integration parameters. The boundary terms in the above action ensure that its associated variational principle features fixation of induced metric and the dilaton on the boundary. Symbolically, we could collect the boundary terms into one expression

$$(\mathcal{S}^I)_{\partial\mathcal{M}} = \frac{1}{2} \alpha \int_{\partial\mathcal{M}} dx \sqrt{|g|} e^{-2\Phi} \mathcal{K} , \quad (3.3)$$

where here $\mathcal{K}$ is used for the extrinsic curvature over the whole boundary $\bar{T}_i \cup \bar{T}_o \cup t' \cup t''$. The corner-point contributions in (3.1) are included because, though the corner points are a set of measure zero in the integration of $e^{-2\Phi} \mathcal{K}$ over all of $\partial\mathcal{M}$, the term $\mathcal{K}$ becomes infinite at the corners, since the boundary normal changes discontinuously from u^a to $\bar{n}^a$ at these points. [10] Finally, we write $\mathcal{S}^I$ for the action, because we anticipate the need to append to the action a Gibbons-Hawking subtraction term $-\mathcal{S}^0[\bar{N}, \Lambda, \Phi]$ (a functional of the fixed boundary data). [12,6] In this case the full action would be $\mathcal{S} \equiv \mathcal{S}^I - \mathcal{S}^0$. We briefly consider the more general action later in an appendix. Also, we could add to the action a matter contribution $\mathcal{S}^m$. Most of our work in this section on quasilocal energy-momentum would be unaffected by an $\mathcal{S}^m$ contribution to the action, as long as the matter fields were minimally coupled. However, an $\mathcal{S}^m$ contribution would affect the following sections devoted to the canonical reduction. Therefore, for the sake of simplicity we do not consider an $\mathcal{S}^m$ further.

We are interested in (vacuum) spherically symmetric general relativity (SSGR). As was first noted by Thome, Isaak, and Hájíček, [13] a suitable action for SSGR is given by (3.1) with the choices $y = 1$, $\alpha = \lambda^{-2}$. Note that for the SSGR case, we set α equal to the *dimensionful* constant λ^{-2} . This gives our action the units of action in four dimensions. An appendix shows that this action can be obtained via a reduction by spherical symmetry of a covariant first-order version [14,10] of the Einstein-Hilbert action (where the *four dimensional* action principle is associated with the time history ${}^4\mathcal{M}$ of a spatial region bounded by concentric spheres). In this correspondence with SSGR, it turns out that the radius of a round sphere is given by

$$R = \sqrt{\frac{A}{4\pi}} = \frac{e^{-\Phi}}{\lambda} , \quad (3.4)$$

where A stands for the proper area of the sphere.

In this paper we also consider (vacuum) two-dimensional dilaton gravity (2dDG), which corresponds to $y = 2$ and α taken to be a dimensionless positive number. Often α is chosen to be $2/\pi$, but α remains essentially arbitrary. The arbitrariness of α for vacuum 2dDG corresponds to the freedom to shift the dilaton by a constant $\Phi \rightarrow \Phi - \log \sqrt{\alpha'}$ (a freedom not present for the SSGR action). Under such a shift, $\alpha \rightarrow \alpha'' = \alpha\alpha'$. We would actually prefer to set $\alpha = 2$ for reasons which become clear later. Adding the right matter contribution

S^m to the vacuum 2dDG action, we would obtain the CGHS model. [9] Though we do not consider this matter contribution, the expressions for quasilocal energy-momentum that we derive in this section are nevertheless valid for the full CGHS theory. For the SSGR case the action $\mathcal{S}^I$ has dimensions of length-squared, while for 2dDG the action $\mathcal{S}^I$ is dimensionless. This difference in units will propagate throughout all the formulae to follow. However, the freedom of allowing α to be either λ^{-2} or a plain number will automatically keep track of the correct units for both cases. For the SSGR case ($y = 1, \alpha = \lambda^{-2}$) all of our conventions have been tailored to match those of KVK and LW.

The first step is to compute the variation of the action. One can compute the variation in a number of ways, but the fastest way is the following. Note that $\mathcal{R}$ is a pure divergence, and then perform an integration by parts on the $\mathcal{R}$ term in the action. This leads to cancelation of most of the boundary terms. This short calculation and resulting form of the action are given later in the discussion in the text preceding (4.3). Vary the resulting form of the action (4.3) to find

$$\delta\mathcal{S}^I = (\text{terms giving the equations of motion}) + \int_{t'}^{t''} dr (P_\Lambda \delta\Lambda + P_\Phi \delta\Phi) + \int_{\bar{\mathcal{T}}} dt \left(\bar{\Pi}_{\bar{N}} \delta\bar{N} + \bar{\Pi}_\Phi \delta\Phi \right) + \alpha e^{-2\Phi} \eta \delta\Phi \Big|_{B'}^{B''}, \quad (3.5)$$

where we have defined the momenta

$$P_\Lambda \equiv \alpha e^{-2\Phi} u^a \nabla_a \Phi \quad (3.6a)$$

$$P_\Phi \equiv -\alpha e^{-2\Phi} \Lambda (\mathcal{K} + y u^a \nabla_a \Phi) \quad (3.6b)$$

$$\bar{\Pi}_{\bar{N}} \equiv -\alpha e^{-2\Phi} \bar{n}^a \nabla_a \Phi \quad (3.6c)$$

$$\bar{\Pi}_\Phi \equiv \alpha e^{-2\Phi} \bar{N} \left(\bar{\nu} + y \bar{n}^a \nabla_a \Phi \right). \quad (3.6d)$$

Inspection of the variation of the action shows that P_Λ is the gravitational momentum conjugate to Λ . Likewise, $\bar{\Pi}_{\bar{N}}$ is the gravitational momentum conjugate to $\bar{N}$, where now conjugacy is defined with respect to the $\bar{\mathcal{T}}$ boundary. The momentum conjugate to the dilaton field is P_Φ . The momentum $\bar{\Pi}_\Phi$ is the $\bar{\mathcal{T}}$ boundary analog of P_Φ . Note that one might try to define a momentum $p_\Phi \equiv \alpha e^{-2\Phi} \eta$ in some sense conjugate to Φ at the corners. However, we chose not to do this. Therefore, strictly speaking, the canonical action we consider later has a mixed Hamiltonian-Lagrangian form. If the corner terms in the action (3.1) had not been included, then the variational principle would have featured fixation of η on the corner points. Fixing Φ at the corners seems to be more in harmony with the fact that Φ is fixed on t', t'' , and the $\bar{\mathcal{T}}$ boundary.

Our momenta P_Λ and P_Φ agree with the analysis of KVK. To see this take the SSGR case and use the fact that $P_R = -(1/R)P_\Phi$. One then finds precisely Kuchař's momenta,

$$P_\Lambda = -N^{-1} R \left(\dot{R} - N^r R' \right) \quad (3.7a)$$

$$P_R = -N^{-1} \left[R \left(\dot{\Lambda} - (\Lambda N^r)' \right) + \Lambda \left(\dot{R} - N^r R' \right) \right]. \quad (3.7b)$$

To get the last expression, we have used the definition of $\mathcal{K}$ given in the preliminary section to find $\mathcal{K} = -(N\Lambda)^{-1} [\dot{\Lambda} - (\Lambda N^r)']$. (Note, however, that due to well-entrenched notation for the dilaton, we unfortunately must break with the KVK convention of only using Greek letters to represent spatial densities like Λ . Though represented by a Greek letter, the dilaton Φ is a scalar and its momentum P_Φ is a density.)

B. Quasilocal energy-momentum

As advertised, our variational principle has been rigged so that the induced metric $(\Lambda^2, -\bar{N}^2)$ and the dilaton Φ are fixed as boundary data. In particular, the lapse of proper time between the initial and final slices is fixed in the variational principle, since this information is encoded in the $\bar{\mathcal{T}}$ boundary metric $-\bar{N}^2$. This is the key feature exploited in the Brown-York theory [6,7] of quasilocal energy-momentum in general relativity. Following the basic line of reasoning in this theory, we “read off” from the variation of the action what geometric expressions play the role of quasilocal energy and momentum in our theory.

We begin by writing the boundary terms in the variation (3.5) of the action in the following suggestive way:

$$(\delta\mathcal{S}^1)_{\partial\mathcal{M}} = - \int_{t'}^{t''} dr (J\delta\Lambda + \Lambda T\delta\Phi) - \int_{\bar{\mathcal{T}}} dt (\bar{E}\delta\bar{N} + \bar{N}\bar{S}\delta\Phi) + \alpha e^{-2\Phi} \eta \delta\Phi \Big|_{B'}^{B''}, \quad (3.8)$$

where we have defined the *scalars*

$$\bar{E} \equiv -\bar{\Pi}_{\bar{N}} = \alpha e^{-2\Phi} \bar{n}^a \nabla_a \Phi \quad (3.9a)$$

$$J \equiv -P_{\Lambda} = -\alpha e^{-2\Phi} u^a \nabla_a \Phi \quad (3.9b)$$

$$\bar{S} \equiv -\bar{\Pi}_{\Phi}/\bar{N} = -\alpha e^{-2\Phi} (\bar{\vartheta} + y \bar{n}^a \nabla_a \Phi) \quad (3.9c)$$

$$T \equiv -P_{\Phi}/\Lambda = \alpha e^{-2\Phi} (\mathcal{K} + y u^a \nabla_a \Phi). \quad (3.9d)$$

We interpret $\bar{E}$ as the *quasilocal energy* associated with the $\bar{u}^a$ observers at the boundary. It is convenient to associate a separate energy with each boundary point of $\bar{\Sigma}$. For instance, $\bar{E}_o \equiv \bar{E}|_{B_o}$ is the energy associated with the outer boundary point. However, properly speaking, the full quasilocal energy associated with the $\bar{\Sigma}$ gravitational and dilaton fields is the sum

$$\bar{E}|_{B_o} + \bar{E}|_{B_i} = \alpha e^{-2\Phi} \bar{n}^a \nabla_a \Phi \Big|_{B_o} - \alpha e^{-2\Phi} \bar{n}^a \nabla_a \Phi \Big|_{B_i} \quad (3.10)$$

of the inner and outer boundary point contributions. Notice that the full energy above is associated with a slice $\bar{\Sigma}$ which has $\bar{u}^a$ as its timelike normal *at the $\bar{\mathcal{T}}$ boundary*.³ Also notice that, when evaluated on solutions to the field equations, the outer-boundary contribution to the energy $\bar{E}_o$, for example, is minus the rate of change of the classical action (or Hamilton-Jacobi principal function) with respect to a unit stretch in $\bar{N}|_{\bar{\mathcal{T}}_o}$, where $\bar{N}|_{\bar{\mathcal{T}}_o}$ controls the lapse of proper time between neighboring points on $\bar{\mathcal{T}}_o$. [6]

The quasilocal energy associated with the u^a observers at the boundary is

$$E \equiv \alpha e^{-2\Phi} n^a \nabla_a \Phi, \quad (3.11)$$

and sum $E_i + E_o$ is the full quasilocal energy for a Σ slice. The energy E depends solely on the Cauchy data of Σ in the same way that $\bar{E}$ depends solely on the Cauchy data of

³Technically, $\bar{E}_i + \bar{E}_o$ is the energy associated with *any* slice in the equivalence class determined by this condition. The slice $\bar{\Sigma}$ can be “wiggled” in the interior as long as its ends remain clamped to the $\bar{\mathcal{T}}$ boundary. Similar comments are relevant for the Σ energy $E_i + E_o$.

$\bar{\Sigma}$. It proves useful in the next section to have an energy expression for *each* point⁴ of Σ . However, we have a sign ambiguity, because each Σ point could be viewed as an inner or an outer boundary point. For the sake of definiteness, define

$$\mathbf{E} \equiv \alpha e^{-2\Phi} \mathbf{n}^a \nabla_a \Phi = \alpha e^{-2\Phi} \Phi' / \Lambda. \quad (3.12)$$

Of course, $E = \pm \mathbf{E}$, depending on whether it is evaluated at the outer or the inner boundary. For some expressions, like E^2 , the sign ambiguity cancels, so we can use E^2 or $\mathbf{E}^2$.

We consider J to be a *quasilocal momentum*. Notice that on-shell $J|_{t''}$, for instance, is minus the rate of change of the Hamilton-Jacobi principal function with respect to a unit stretch in $\Lambda|_{t''}$, where $\Lambda|_{t''}$ controls the lapse of proper distance between neighboring radial leaves B of t'' . Such a variation in the boundary data corresponds to a dilation of t'' . Again, we associate a J with each point of a generic Σ slice. At a glance, one sees that $J = -P_\Lambda$ depends solely on Σ Cauchy data.

C. Boost relations and invariants

The Eulerian observers of $\bar{\Sigma}$ at the $\bar{\mathcal{T}}$ boundary coincide with the natural observers in the boundary. We may define a set

$$\bar{E} = \alpha e^{-2\Phi} \bar{n}^a \nabla_a \Phi \quad (3.13a)$$

$$\bar{J} = -\alpha e^{-2\Phi} \bar{u}^a \nabla_a \Phi \quad (3.13b)$$

$$\bar{S} = -\alpha e^{-2\Phi} (\bar{\vartheta} + y \bar{n}^a \nabla_a \Phi) \quad (3.13c)$$

$$\bar{T} = \alpha e^{-2\Phi} (\bar{\mathcal{K}} + y \bar{u}^a \nabla_a \Phi) \quad (3.13d)$$

of quasilocal quantities for such observers. Now the slice Σ need not be clamped to the $\bar{\mathcal{T}}$ boundary in our formalism. Hence, in general Σ and $\bar{\Sigma}$ are different slices which intersect at the same boundary point of interest. We will refer to a switch of the spatial slice spanning a particular boundary point as a *generalized boost* or simply a *boost*. Properly speaking, a generalized boost is a switch of the equivalence class of spanning slices. The behavior

$$\bar{E} = \gamma E - v \gamma J \quad (3.14a)$$

$$\bar{J} = \gamma J - v \gamma E \quad (3.14b)$$

$$\bar{S} = \gamma S - v \gamma T + \alpha e^{-2\Phi} \bar{u}^a \nabla_a \eta \quad (3.14c)$$

$$\bar{T} = \gamma T - v \gamma S - \alpha e^{-2\Phi} \bar{n}^a \nabla_a \eta \quad (3.14d)$$

of the quasilocal quantities under boosts follows immediately from the boost relations (2.6) and the splittings (2.7). In (3.14c,d) the unbarred version of $\bar{S}$ is

$$S \equiv -\alpha e^{-2\Phi} \left(-a_b n^b + y n^b \nabla_b \Phi \right). \quad (3.15)$$

⁴Recall that the Σ points are spheres, at least in the SSGR case. So we are not really defining a local energy, and certainly not a local energy *density*.

Notice that E , J , S , and T have the same dependence on Σ Cauchy data that $\bar{E}$, $\bar{J}$, $\bar{S}$ and $\bar{T}$ have on $\bar{\Sigma}$ Cauchy data. (However, due to the appearance of acceleration terms, both S and $\bar{S}$ do not depend solely on Cauchy data.)

Clearly the expression $-E^2 + J^2$ is invariant under boosts. We may multiply it by any function of the dilaton field or add to it any function of the dilaton field and retain a boost-invariant expression. Therefore, it is not completely unnatural to introduce the invariant

$$M = (2\alpha\lambda)^{-1} e^{y\Phi} \left[-E^2 + J^2 + (\alpha\lambda)^2 \exp[-2y\Phi] \right]. \quad (3.16)$$

Note that M has units of length for the SSGR case and units of inverse length for 2dDG. It turns out that on-shell (on solutions to the field equations) M is a completely conserved quantity (constant in time and space). Moreover, for the case of SSGR we find that $M = M_s$, where M_s is Kuchař's canonical expression for the Schwarzschild mass parameter,

$$M_s = \frac{P_\Lambda^2}{2R} - \frac{R}{2} \left(\frac{R'}{\Lambda} \right)^2 + \frac{R}{2}. \quad (3.17)$$

From the four-dimensional spacetime perspective, the expression for M_s corresponds to several mass definitions in general relativity, when spherical symmetry is assumed. One is the Hawking mass [16,17]

$$M_H = \frac{1}{8\pi} \sqrt{\frac{A}{16\pi}} \int_B d^2x \sqrt{\sigma} \left[-\frac{1}{2}(k^2 - \ell^2) + {}^2R \right], \quad (3.18)$$

where 2R is the intrinsic scalar curvature of the two-surface B (in our case a round sphere), and k is the trace of the intrinsic curvature of B as embedded in a spanning (three-dimensional) Σ slice. Likewise, ℓ is the trace of the extrinsic curvature of B as embedded in the timelike three-surface generated by the integral curves of the Σ normal, denoted by u^μ for four-dimensions. The boost-invariant combination $\frac{1}{2}(k^2 - \ell^2)$ is more often written as a product of the expansions associated with the ingoing and outgoing null normals to B , and the factor A , the area of B , ensures that the whole expression has units of energy. For Schwarzschild M_H equals the mass parameter of the solution even for finite two-spheres. For general closed two-surfaces in general spacetimes, the Hawking expression can be “built” as a combination of “quasilocal boost invariants.” [7,15]

For the 2dDG case set $M_W = 2M/\alpha$. By expressing M_W in covariant form,

$$M_W = \lambda^{-1} e^{-2\Phi} \left(\lambda^2 - g^{ab} \nabla_a \Phi \nabla_b \Phi \right), \quad (3.19)$$

we see that it is the “local mass” of Tada and Uehara. [18] Such a quantity was also considered by Frolov in Ref. [19]. With an argument originally given by KVK for the Schwarzschild case, the appendix shows that M_W is the canonical expression for the mass parameter of the Witten black hole. The appendix also shows that the ADM energy [2] at spatial infinity (associated with the preferred static-time slices) of the Witten black hole is the on-shell value of M . This is the reason we would prefer to set $\alpha = 2$.

IV. CANONICAL THEORY

This section is devoted to the canonical form of the theory. We first sketch the Legendre transformation which yields the canonical form of the action. We then vary the canonical action, paying strict attention to all boundary terms. Finally, we consider the canonical transformation of KVK and write down a new-canonical-variable version of the action (3.1) which is particularly amenable to canonical reduction.

A. Form of the canonical action

Passage to the canonical form of the action (3.1) demands that we write the action in $1+1$ form as a preliminary step. This is easily done with three ingredients. The first is the splitting result (2.7b) for the $\bar{T}$ boundary extrinsic curvature $\bar{\vartheta}$. The second is the identity

$$2yg^{ab}\nabla_a\Phi\nabla_b\Phi = 2y\left[-\left(\frac{\dot{\Phi}}{N}\right)^2 + 2N^r\left(\frac{\dot{\Phi}\Phi'}{N^2}\right) + \left(\frac{\Phi'}{\Lambda\gamma}\right)^2\right], \quad (4.1)$$

where γ is the relativistic factor of the preliminary section. The third and final ingredient is the realization that for our two dimensional spacetime the Ricci scalar is a pure divergence,

$$\mathcal{R} = -2\nabla_a(\mathcal{K}u^a + a_b n^b n^a). \quad (4.2)$$

Using these three ingredients, one can quickly cast (3.1) into the form

$$\mathcal{S}^I = \frac{1}{4}\alpha\int_{\mathcal{M}} d^2x N\Lambda e^{-2\Phi}X + \int_{\bar{T}} dt \alpha e^{-2\Phi}\eta\dot{\Phi}, \quad (4.3)$$

where X is the expression

$$\begin{aligned} X = & -\frac{4}{N\Lambda}\left[\Lambda\mathcal{K}\left(\dot{\Phi} - N^r\Phi'\right) + \frac{N'\Phi'}{\Lambda}\right] \\ & + 2y\left[-\left(\frac{\dot{\Phi}}{N}\right)^2 + 2N^r\left(\frac{\dot{\Phi}\Phi'}{N^2}\right) + \left(\frac{\Phi'}{\Lambda\gamma}\right)^2\right] + 2\lambda^2 y \exp[(2-y)2\Phi]. \end{aligned} \quad (4.4)$$

Performing the usual calculations with (4.3), one finds the following canonical action:

$$\mathcal{S}^I = \int_{\mathcal{M}} d^2x \left(P_{\Lambda}\dot{\Lambda} + P_{\Phi}\dot{\Phi} - N\mathcal{H} - N^r\mathcal{H}_r\right) + \int_{\bar{T}} dt \left(\alpha e^{-2\Phi}\eta\dot{\Phi} - \bar{N}\bar{E}\right), \quad (4.5)$$

where here $\bar{E}$ is short-hand for $\gamma E - v\gamma J$. It is important to realize that in the canonical picture the equation $\bar{E} = \alpha e^{-2\Phi}\bar{n}^a\nabla_a\Phi$ is not necessarily valid, for equality implicitly assumes the *canonical equation of motion* $P_{\Lambda} = \alpha e^{-2\Phi}u^a\nabla_a\Phi$. Respectively, the Hamiltonian constraint and the momentum constraint have the form

$$\begin{aligned} \mathcal{H} = & \alpha^{-1}e^{2\Phi}\left[P_{\Phi}P_{\Lambda} + \frac{1}{2}y\Lambda(P_{\Lambda})^2\right] \\ & + \alpha e^{-2\Phi}\left[\left(2 - \frac{1}{2}y\right)\frac{\Phi'^2}{\Lambda} - \frac{\Phi''}{\Lambda} + \frac{\Phi'\Lambda'}{\Lambda^2} - \frac{1}{2}\lambda^2\Lambda y \exp[(2-y)2\Phi]\right] \end{aligned} \quad (4.6a)$$

$$\mathcal{H}_r = P_{\Phi}\Phi' - \Lambda P'_{\Lambda}. \quad (4.6b)$$

Notice that it is P_{Λ} which appears differentiated in the momentum constraint $\mathcal{H}_r$. This is to be expected, as Λ is a scalar density.

B. Variation of the canonical action

Straightforward manipulations establish that the variation of the canonical action is

$$\begin{aligned} \delta \mathcal{S}^I = & \text{(terms which enforce the constraints and give the} \\ & \text{Hamiltonian equations of motion)} + \int_{t'}^{t''} dr (P_\Lambda \delta \Lambda + P_\Phi \delta \Phi) \\ & - \int_{\bar{T}} dt \left[\bar{E} \delta \bar{N} + \bar{N} \bar{S} \delta \Phi - \left(\bar{N} \bar{J} + \alpha e^{-2\Phi} \dot{\Phi} \right) \delta \eta \right] + \alpha e^{-2\Phi} \eta \delta \Phi \Big|_{B'}^{B''}, \end{aligned} \quad (4.7)$$

where here $\bar{E}$, $\bar{J}$, and $\bar{S}$ are short-hand for the expressions

$$\bar{E} = \gamma E - v \gamma J \quad (4.8a)$$

$$\bar{J} = \gamma J - v \gamma E \quad (4.8b)$$

$$\bar{S} = \gamma S - v \gamma T + \alpha e^{-2\Phi} \bar{u}^a \nabla_a \eta. \quad (4.8c)$$

As mentioned, one should be careful, for while E , J , S , and T are built from the canonical variables $(\Lambda, P_\Lambda; \Phi, P_\Phi)$ in the same way as before (but S does not depend solely on the canonical variables), *now the momenta P_Λ and P_Φ need not have the forms given in (3.6)* (which are canonical equations of motion). The term which appears multiplied by the variation $\delta \eta$ vanishes when the canonical equation of motion for P_Λ holds. Therefore, η is not a quantity which is held fixed in our variational principle.

C. Canonical Transformation

In this subsection we perform Kuchař's canonical transformation on the phase-space pairs $(\Lambda, P_\Lambda; \Phi, P_\Phi)$. In order to grasp the underlying hyperbolic geometry of this canonical transformation, we first need to collect a few results and observations.

Consider a black-hole solution which extremizes the action (3.1) (either a Schwarzschild black hole or a Witten black hole, depending on whether y is 1 or 2). Associated with this solution, there is a preferred family of static-time slices, the collection of constant-Killing-time level surfaces. For the Schwarzschild-black-hole case let $T(t, r)$ denote the Killing time, and for the Witten-black-hole case let $\tau(t, r)$ denote the Killing time. Now, given a particular $\mathcal{M}$ point B , we may interpret it as a boundary point of the static-time slice $\tilde{\Sigma}$ which contains it. Our construction defines the *rest frame* $(\tilde{u}^a, \tilde{n}^a)$ at B , where $\tilde{u}^a$ is the normal of $\tilde{\Sigma}$ as embedded in $\mathcal{M}$ and $\tilde{n}^a$ is the normal of B as embedded in $\tilde{\Sigma}$. If B is also considered to be a point of the $\bar{T}$ boundary, then in general $\tilde{\Sigma}$ does not define the same frame at B as the slice Σ or the slice $\bar{\Sigma}$ considered before. We know how to compute the energy $\tilde{E}$ and momentum $\tilde{J}$ of the bounded static-time slice $\tilde{\Sigma}$,

$$\begin{aligned} \tilde{E} &= \alpha e^{-2\Phi} \tilde{n}^a \nabla_a \Phi \\ \tilde{J} &= -\alpha e^{-2\Phi} \tilde{u}^a \nabla_a \Phi. \end{aligned} \quad (4.8)$$

Clearly $\tilde{E}$ and $\tilde{J}$ depend on the Cauchy data of $\tilde{\Sigma}$ in the same way that E and J depend on the Cauchy data of Σ .

Now, it is a fact that $\tilde{J} = 0$, which is why we refer to $(\tilde{u}^a, \tilde{n}^a)$ as the rest frame at B . The existence of the rest frame at B leads to our key observation: *at B the parameter φ associated with the boost from the frame (u^a, n^a) defined by Σ to the rest frame $(\tilde{u}^a, \tilde{n}^a)$ is determined by the canonical variables of Σ .*⁵ Indeed, with $w \equiv J/E$ the boost from the Σ frame to the rest frame is parameterized by

$$\varphi \equiv \frac{1}{2} \log \left| \frac{1+w}{1-w} \right| = \frac{1}{2} \log \left| \frac{E+J}{E-J} \right|. \quad (4.9)$$

Notice that $\tilde{u}^a = \psi u^a + w \psi n^a$ with $\psi = (1-w^2)^{-1/2}$. We know the expression (4.9) is correct, because our general theory of boosted quasilocal energy-momentum implies that $E = \psi \tilde{E}$ and $J = w \psi \tilde{E}$. To get these last relations, assume that the $\tilde{T}$ boundary is generated by the $\tilde{u}^a$ Eulerian history of B points (identify “tildes” with “bars” and w with v) and use (3.14a,b). At this stage we have a sign ambiguity in our expressions, since we did not say whether B is an inner or an outer boundary point. For the sake of definiteness in what follows, we often want to assume that B is taken as an outer boundary point and make this distinction notationally. Therefore, we use $\mathbf{E}$, defined before in (3.12), and also $\mathbf{w} = J/\mathbf{E}$ which defines $\varphi \equiv \frac{1}{2} \log |(1+\mathbf{w})/(1-\mathbf{w})|$. Notice that the sign ambiguities cancel in the expression $E\varphi = \mathbf{E}\varphi$. One should be careful when dealing with $\mathbf{w}$, ψ , and φ . Since for a classical solution the dilaton is a “bad” radial coordinate in the sense that $\Phi' < 0$, our *unreferenced* energy $\mathbf{E}$ is negative. The (in our case positive) expression for the quasilocal energy with the flat-space reference contribution is considered in the appendix and at length in Ref. [6]. Therefore, since square roots are by convention positive, $\psi = -\mathbf{E}/\sqrt{E^2 - J^2}$.

With our canonical expression for φ we straightforwardly write down a new set of constraints which generate proper unit displacements with respect to the static-time slices,

$$\mathbf{H} = \psi \mathcal{H} + \mathbf{w} \psi (\mathcal{H}_r / \Lambda) \quad (4.10a)$$

$$\mathbf{H}_\perp = \psi (\mathcal{H}_r / \Lambda) + \mathbf{w} \psi \mathcal{H}. \quad (4.10b)$$

Since these new constraints depend only on the canonical variables of Σ , we can consider them off the constraint surface in phase space. However, when computing a Poisson bracket $\{G, \mathbf{H}\}$, where G is a functional of the canonical variables, one finds that all of the “dangerous” brackets $\{G, \mathbf{w}\}$ which arise come multiplied by either by factor of $\mathcal{H}$ or $\mathcal{H}_r$. Therefore, on-shell $\mathbf{H}$ generates proper unit displacement of the Σ slice in the $\tilde{u}^a$ direction. Likewise, $\mathbf{H}_\perp$ generates proper unit displacement of the Σ slice in the $\tilde{n}^a$ direction. As we show below, these generators are closely related to the momenta P_T and P_R considered in KVK.

Define the new canonical variables in terms of the old ones as follows:

$$M = \frac{1}{2} \alpha \lambda e^{-y\Phi} (1 - F) \quad (4.11a)$$

$$P_M = (\alpha \lambda)^{-1} e^{y\Phi} F^{-1} \Lambda P_\Lambda \quad (4.11b)$$

$$\Psi = \Phi \quad (4.11c)$$

$$P_\Psi = P_\Phi + \frac{1}{2} y (1 + F^{-1}) \Lambda P_\Lambda + \alpha e^{-2\Phi} \varphi', \quad (4.11d)$$

⁵Which need to be known only in a tiny neighborhood of B .

where M is the boost invariant (3.16) and in terms of the old variables F and φ are short-hand for

$$F = (\alpha\lambda)^{-2} \exp[(y-2)2\Phi] \left[(\alpha\Phi'/\Lambda)^2 - (e^{2\Phi}P_\Lambda)^2 \right] \quad (4.12a)$$

$$\varphi = \frac{1}{2} \log \left| \frac{\alpha e^{-2\Phi}\Phi' - \Lambda P_\Lambda}{\alpha e^{-2\Phi}\Phi' + \Lambda P_\Lambda} \right|. \quad (4.12b)$$

Notice that $\varphi = \frac{1}{2} \log |F_-/F_+|$, where

$$F_\pm = (\alpha\lambda)^{-1} e^{y\Phi} (\mathbf{E} \mp J). \quad (4.13)$$

Evidently then, another expression for F of key importance is

$$F = F_+ F_- = (\alpha\lambda)^{-2} e^{y2\Phi} (E^2 - J^2). \quad (4.14)$$

For the SSGR case $e^{-\Psi} = \lambda R$ and $P_\Psi = -R P_R$, in the notation of KVK and LW. Also for this case, one finds that $P_M = -T'$, because as shown in KVK the canonical expression for (minus) the radial derivative of the Killing time is given by $-T' = (RF)^{-1} \Lambda P_\Lambda$. The situation is the same for 2dDG, as in this case $P_M = -\tau'$. We show in the appendix that the canonical expression for (minus) the radial derivative of the Witten-black-hole Killing time is $-\tau' = (\alpha\lambda)^{-1} e^{2\Phi} F^{-1} \Lambda P_\Lambda$. One may prove that the transformation $(\Lambda, P_\Lambda; \Phi, P_\Phi) \rightarrow (M, P_M; \Psi, P_\Psi)$ is canonical for our boundary conditions by verifying the identity

$$P_\Lambda \delta \Lambda + P_\Phi \delta \Phi - P_M \delta M - P_\Psi \delta \Psi = \delta [\Lambda(-J + E\varphi)] - (\alpha e^{-2\Phi} \delta \Phi \varphi)', \quad (4.15)$$

which upon integration over r shows that the difference between the old Liouville form and the new Liouville is an exact form. Hence, the transformation is canonical.

Recall that for simplicity we wish to restrict our attention to the right-static region of the relevant Kruskal diagram. Now, in fact, for both SSGR and 2dDG F is the canonical expression for $\tilde{N}^2$, where $\tilde{N}$ is the lapse function associated with the static-time slices. On-shell, the event horizon of a particular black-hole solution is the locus of points determined by $F = 0$. We may ensure that we are working exclusively in a static region of the Kruskal diagram by choosing our boundary conditions appropriately and by excluding solutions for which $F = 0$ somewhere on $\mathcal{M}$. Where F is nonzero, the above transformation may be inverted,

$$\Lambda = \left[\lambda^{-2} F^{-1} (\Psi')^2 \exp[(y-2)2\Psi] - F(P_M)^2 \right]^{1/2} \quad (4.16a)$$

$$P_\Lambda = \frac{\alpha \lambda e^{-y\Psi} F P_M}{[\lambda^{-2} F^{-1} (\Psi')^2 \exp[(y-2)2\Psi] - F(P_M)^2]^{1/2}} \quad (4.16b)$$

$$\Phi = \Psi \quad (4.16c)$$

$$P_\Phi = P_\Psi - \frac{1}{2} \alpha \lambda y e^{-y\Psi} P_M (1 + F) - \alpha e^{-2\Psi} \varphi', \quad (4.16d)$$

where in terms of the new variables F and φ are short-hand for

$$F = 1 - 2(\alpha\lambda)^{-1}e^{y\Psi}M \quad (4.17a)$$

$$\varphi = \frac{1}{2} \log \left| \frac{\Psi' - \lambda \exp[(2-y)\Psi] F P_M}{\Psi' + \lambda \exp[(2-y)\Psi] F P_M} \right|. \quad (4.17b)$$

As mentioned in the introduction, KVK considers the canonical transformation and its inverse in all regions of the Kruskal diagram for SSGR. Moreover, this reference provides a detailed treatment of the (singular) behavior of the transformation at the horizon. We expect that a similar treatment with essentially the same results can be carried out for the 2dDG case.

As shown in KVK for the SSGR case, the payoff obtained by using the new variables comes when considering the constraints (4.6). Since on solutions to the constraints M is a constant, one knows that M' must be a sum of constraints. Indeed, direct calculation establishes that

$$M' = -F^{1/2}\mathbf{H}. \quad (4.18)$$

Since $F^{1/2}$ is the lapse $\tilde{N}$, we see that $-M'$ is the generator of Killing-time evolution. For this reason, KVK uses the notation $P_T = -M'$ (and we would likewise set $P_\tau = -M'$ for the 2dDG case). Furthermore,

$$P_\Psi = -\lambda^{-1} \exp[(y-2)\Phi] F^{-1/2} \mathbf{H}_- \quad (4.19)$$

is also a sum of constraints and so weakly vanishes. KVK and LW set $P_R = F^{-1/2} \mathbf{H}_-$. As described in KVK, the momenta P_T and P_R respectively generate coordinate-scaled displacements of the Σ slice along lines of constant $R = R$ and T (or τ for 2dDG) in the Kruskal diagram. Kuchař obtains these interpretations for P_T and P_R via a “parameterization at infinity” and canonical transformations subsequent to the main one (4.11). We have obtained the exact same interpretations via a different route.

The new canonical variables are related to the new constraints (4.10) in a very simply way: $\mathbf{H} = -F^{-1/2}M'$ and $\mathbf{H}_- = -\lambda \exp[(2-y)\Psi] F^{1/2} P_\Psi$. Using these relations, we may write the old constraints in terms of the new variables as

$$\begin{aligned} \mathcal{H} &= \psi \mathbf{H} - \mathbf{w} \psi \mathbf{H}_- \\ \mathcal{H}_r &= \Lambda (\psi \mathbf{H}_- - \mathbf{w} \psi \mathbf{H}), \end{aligned} \quad (4.20)$$

where here Λ is given in (4.16a) and $\mathbf{w}$ must be expressed in terms of the new variables. With the list (4.16), it is not hard to show that

$$\mathbf{w} = -\lambda(\Psi')^{-1} \exp[(2-y)\Psi] F P_M. \quad (4.21)$$

It is now straightforward, if tedious, to express the old constraints in terms of the new variables,

$$\mathcal{H} = \frac{\lambda^{-1} F^{-1} \Psi' \exp[(y-2)\Psi] M' + \lambda F \exp[(2-y)\Psi] P_\Psi P_M}{[\lambda^{-2} F^{-1} (\Psi')^2 \exp[(y-2)2\Psi] - F(P_M)^2]^{1/2}} \quad (4.22)$$

$$\mathcal{H}_r = P_M M' + P_\Psi \Psi'.$$

As noted in KVK, in terms of the new variables it is relatively simple to show that the Poisson bracket of M with either $\mathcal{H}$ or $\mathcal{H}_r$ vanishes weakly.

D. Canonical reduction

The goal of this subsection is to use the new canonical variables to find a reduced action principle which -in a certain sense- corresponds to the canonical action (4.5). However, we will be adding boundary terms to (4.5) before the reduction is made, so we should clearly state what we have in mind to begin with and why. Several aspects of the path-integral formulation of gravitational thermodynamics motivate fixation of $\bar{N}$ and Φ on the $\bar{\mathcal{T}}$ boundary as the features of central importance which need to be preserved as we modify the original action (4.5). In path-integral expressions for gravitational partition functions, the sum over histories includes only spacetimes for which the initial and final slices are identified. In this scenario the gauge-invariant information of $\bar{N}$ (the lapse of proper time between the identified initial slice t' and final slice t'') is essentially the inverse temperature, which is fixed in the canonical ensemble. [5] Regarding the fixation of the dilaton Φ on the $\bar{\mathcal{T}}$ boundary, from a four-dimensional perspective this feature allows the area of the boundary of the system to be fixed as a boundary condition.

In what follows we modify our original canonical action (4.5) in several steps. Each step is -at least heuristically- justified. The result will be an action $\mathcal{S}_\ddagger^I$ (for the SSGR case this action is closely related to one considered by LW) which is particularly amenable to canonical reduction via the new variables. Moreover, as we will explicitly demonstrate, the new action $\mathcal{S}_\ddagger^I$ retains fixation of $\bar{N}$ and Φ on the $\bar{\mathcal{T}}$ boundary, important for the above mentioned reasons, as features of its associated variational principle. Our analysis provides some conceptual justification for several technical steps taken in LW.

Let us go through the steps of modifying $\mathcal{S}^I$. We know from (4.15) that addition of the boundary terms

$$-\omega[\Lambda, P_\Lambda, \Phi]|_{t'}^{t''} = \int_{t'}^{t''} dr \Lambda (J - E\varphi) \quad (4.23)$$

to the canonical action (4.5) gives the new action

$$\mathcal{S}_\ddagger^I \equiv \int_{\mathcal{M}} d^2x \left(P_M \dot{M} + P_\Psi \dot{\Psi} - N\mathcal{H} - N^r \mathcal{H}_r \right) + \int_{\bar{\mathcal{T}}} dt \left(\alpha e^{-2\Psi} \eta \dot{\Psi} - \bar{N} \bar{E} \right), \quad (4.24)$$

where here we consider all the quantities as expressed in terms of the old variables. The vanishing of the original set of constraints is equivalent to the vanishing of M' and P_Ψ . To take advantage of this fact, re-express the constraint terms in the action as $N\mathcal{H} + N^r \mathcal{H}_r = N^M M' + N^\Psi P_\Psi$, where the new Lagrange multipliers are

$$N^M = -F^{-1/2} (\psi N - \mathbf{w} \psi \Lambda N^r) \quad (4.25a)$$

$$N^\Psi = -\lambda \exp[(2 - y)\Psi] F^{1/2} (\psi \Lambda N^r - \mathbf{w} \psi N). \quad (4.25b)$$

At this stage the new Lagrange multipliers still depend on the old multipliers and the old canonical variables. In particular, note that in terms of the old variables

$$N^\Psi = -\alpha^{-1} e^{2\Phi} \bar{N} (\gamma J - v \gamma \mathbf{E}), \quad (4.26)$$

where we have used the expression (4.14) and the fact that $\mathbf{w} = J/\mathbf{E}$. With the canonical equation of motion for P_Λ , one can show that $N^\Psi = -\alpha^{-1} e^{2\Phi} \bar{N} \bar{J} = \bar{\Phi}$.

Had we merely passed to the new canonical variables, without redefining the Lagrange multipliers, the variational principle associated with (4.24) would have featured fixation of $\bar{N}$ and $\bar{\Phi}$ on the $\bar{\mathcal{T}}$ boundary automatically. However, with the Lagrange multipliers redefined (and in a way which absorbs some of the canonical variables) all bets are off. We must cleverly choose the $\bar{\mathcal{T}}$ boundary term appropriate for the new-variable version of the action. In passing from the old constraints to M' and P_Ψ , we are effectively performing the Lorentz boost from the frame $(u^a, \mathbf{n}^a)$ to the rest frame $(\tilde{u}^a, \tilde{\mathbf{n}}^a)$ at each point on Σ . A point-wise boost⁶ has been performed on the old constraints and Lagrange multipliers. However, we have not included the boundary term in the boost. It seems that the correct way to incorporate the effect of the boost into the boundary term is to reference the existing boost parameter η against the parameter φ associated with the boost to the rest frame. This is achieved by adding a $\bar{\mathcal{T}}$ boundary term to the action $\mathcal{S}_\dagger^I$, with the result

$$\mathcal{S}_\dagger^I \equiv \int_{\mathcal{M}} d^2x \left(P_M \dot{M} + P_\Psi \dot{\Psi} - N^M M' - N^\Psi P_\Psi \right) + \int_{\bar{\mathcal{T}}} dt \left(\alpha e^{-2\Psi} \rho \dot{\Psi} - \bar{N} \bar{E} \right), \quad (4.27)$$

On-shell, the boost parameter in the new action

$$\rho \equiv \eta - \varphi = -\frac{1}{2} \log \left| \frac{\bar{E} + \bar{J}}{\bar{E} - \bar{J}} \right| \quad (4.28)$$

is associated with the local boost between the rest frame $(\tilde{u}^a, \tilde{\mathbf{n}}^a)$ and the boundary frame $(\bar{u}^a, \bar{\mathbf{n}}^a)$. At this stage, the terms $\bar{E}$ and $\bar{J}$ are still short-hand expressions for $\gamma E - v\gamma J$ and $\gamma J - v\gamma E$, respectively.

We now wish to express all terms in the action $\mathcal{S}_\dagger^I$ solely in terms of the new variables and freely variable N^M and N^Ψ . Using the expressions (4.16), one can easily express the quasilocal energy and momentum in terms of the new variables,

$$E = \frac{\epsilon \alpha e^{-2\Psi} \Psi'}{[\lambda^{-2} F^{-1}(\Psi')^2 \exp[(y-2)2\Psi] - F(P_M)^2]^{1/2}} \quad (4.29a)$$

$$J = \frac{-\alpha \lambda e^{-y\Psi} F P_M}{[\lambda^{-2} F^{-1}(\Psi')^2 \exp[(y-2)2\Psi] - F(P_M)^2]^{1/2}}, \quad (4.29b)$$

where the factor $\epsilon \equiv (\bar{n}_a \bar{\mathbf{n}}^a)$ takes care of the appropriate sign on each of the boundary elements. Moreover, we must now regard N and N^r (which along with Λ are hidden in the v 's and γ 's which are in turn hidden in ρ) as depending on the new variables. It is not difficult to invert the relations (4.25) to get the needed expressions. Also, a short calculation shows that the boundary lapse $\bar{N} = N/\gamma$ has a fairly nice expression in terms of the new variables,

$$\bar{N}^2 = F \left(N^M \right)^2 - \lambda^{-2} \exp[(y-2)2\Psi] F^{-1} \left(N^\Psi \right)^2. \quad (4.30)$$

However, the situation at hand remains quite problematic. We would like to use the fact that $M' = 0$ to define a radially independent $\mathbf{m}(t) \equiv M(t)$ with conjugate momentum

⁶Or, depending on the viewpoint, a sphere-wise boost.

$\mathbf{p}(t) \equiv \int_{\Sigma} dr P_M(t, r)$. Since we have seen that P_M is minus the radial derivative of the Killing time, the momentum $\mathbf{p}$ would then be the difference between the Killing-time values for the boundary points B_i and B_o . Indeed, our canonical-reduction goal is to insert the solutions of the constraints into the action to find a reduced action which is expressed in terms of the pair $(\mathbf{m}, \mathbf{p})$. However, P_M appears in the $\bar{\mathcal{T}}$ boundary terms explicitly. Therefore, even if we perform the r integration in the action to define $\mathbf{p}$, the boundary terms still contain factors of P_M .⁷

A solution to the problem at hand is to make an appeal to the equations of motion. For the moment, let us go back to considering the action $\mathcal{S}_{\ddagger}^I$ as depending on the old variables. Notice that *using the canonical equation of motion for P_{Λ}* , one may write

$$\bar{J} = -\alpha e^{-2\Phi}(\dot{\Phi}/\bar{N}) \quad (4.31a)$$

$$\bar{E} = -\epsilon \alpha e^{-2\Phi} \sqrt{\lambda^2 \exp[(2-y)2\Phi] F + (\dot{\Phi}/\bar{N})^2}, \quad (4.31b)$$

where we have appealed to the form (4.14) for F and again used ϵ to take care of the appropriate sign for each boundary element. Again, note that by convention we take the positive square root in (4.31b) and $\bar{E}$ is negative. It is trivial to write these new expressions for $\bar{E}$ and $\bar{J}$ in terms of the new variables. Fortunately, these expressions have no dependence on P_M . Using these expressions instead of those in (4.29), we find that in terms of the new variables our action is again (4.27); but now with (i) $\bar{N}\bar{E} = -\epsilon \alpha e^{-2\Psi} \sqrt{\lambda^2 \exp[(2-y)2\Psi] \bar{N}^2 F + (\dot{\Psi})^2}$, (ii) $\bar{N}^2$ short-hand for the expression (4.30), (iii) F short-hand for expression (4.17a), and (iv) the parameter specifying the boost between the boundary and rest frames given by

$$\rho = -\frac{1}{2} \log \left| \frac{\sqrt{\lambda^2 \exp[(2-y)2\Psi] \bar{N}^2 F + (\dot{\Psi})^2} + \epsilon \dot{\Psi}}{\sqrt{\lambda^2 \exp[(2-y)2\Psi] \bar{N}^2 F + (\dot{\Psi})^2} - \epsilon \dot{\Psi}} \right|. \quad (4.32)$$

Now let us verify that the variational principle associated with (4.27) and the just-given list (i)-(iv) does in fact possess the features that we demand. The constraints and equations of motion are

$$M' = 0 \quad (4.33a)$$

$$P_{\Psi} = 0 \quad (4.33b)$$

$$\dot{M} = 0 \quad (4.33c)$$

$$\dot{P}_M = N^M \quad (4.33d)$$

$$\dot{\Psi} = N^{\Psi} \quad (4.33e)$$

$$\dot{P}_{\Psi} = 0. \quad (4.33f)$$

⁷The reader might suspect that this problem was introduced when we performed the heuristically justified reference $\eta \rightarrow \rho = (\eta - \varphi)$ to get the action $\mathcal{S}_{\ddagger}^I$. But the boundary term in the action $\mathcal{S}_{\ddagger}^I$ suffers from the same problem. Indeed, η is built from v which is in turn built from Λ , and hence in term of the new variables η depends on P_M .

In terms of the old variables, we have already seen that the equation (4.33e) holds when the canonical equation of motion for P_Λ is assumed. Upon variation of the action, we find the boundary terms

$$\begin{aligned} (\delta \mathcal{S}_\dagger^I)_{\partial \mathcal{M}} = \\ \int_{t'}^{t''} dr (P_M \delta M + P_\Psi \delta \Psi) + \int_{\bar{\mathcal{T}}} dt (\bar{\Pi}_{\bar{N}} \delta \bar{N} + \bar{\Pi}_\Psi \delta \Psi + \bar{\Pi}_M \delta M) + \alpha e^{-2\Psi} \rho \delta \Psi \Big|_{B'}^{B''}, \end{aligned} \quad (4.34)$$

where now the $\bar{\mathcal{T}}$ boundary momenta are the following:

$$- \bar{E} = \bar{\Pi}_{\bar{N}} = \epsilon \alpha e^{-2\Psi} \sqrt{\lambda^2 \exp[(2-y)2\Psi] F + (\dot{\Psi}/\bar{N})^2} \quad (4.35a)$$

$$\bar{\Pi}_\Psi = \frac{1}{2} \bar{N} \left[y \bar{E} (1 + F^{-1}) - 2 \alpha e^{-2\Psi} \bar{u}^a \nabla_a \rho \right] \quad (4.35b)$$

$$\bar{\Pi}_M = -\epsilon N^M + (\alpha \lambda)^{-1} e^{y\Psi} F^{-1} \bar{N} \bar{E}. \quad (4.35c)$$

Note that $\bar{\Pi}_\Psi$ is not the same as the $\bar{\Pi}_\Phi$ in the list (3.6). Although the $\bar{\Pi}_{\bar{N}}$ found here is not the expression (3.6c), it agrees with this expression on-shell. Recalling that $\bar{N}^2$ stands for (4.30), one finds that $\bar{\Pi}_M$ vanishes when the equation of motion (4.33e) holds. Therefore, M need not be held fixed on the $\bar{\mathcal{T}}$ boundary in the variational principle associated with $\mathcal{S}_\dagger^I$, as the equations of motion ensure that the boundary terms with $\bar{\Pi}_M$ vanishes for arbitrary variations δM about a classical solution.

The reduced action $I_\dagger^I$, expressed in terms of $\mathbf{m}$ and $\mathbf{p}$ defined earlier, is obtained by solving the constraints and inserting these solutions back into the action $\mathcal{S}_\dagger$. From the result (4.34) for the variation of the action $\mathcal{S}_\dagger^I$, we know that the reduced action,

$$\begin{aligned} I_\dagger^I = \int_{t'}^{t''} dt \mathbf{p} \dot{\mathbf{m}} + \int_{\bar{\mathcal{T}}} dt \alpha e^{-2\Psi} \left[\epsilon \sqrt{\lambda^2 \exp[(2-y)2\Psi] \bar{N}^2 F + (\dot{\Psi})^2} \right. \\ \left. - \frac{1}{2} \dot{\Psi} \log \left| \frac{\sqrt{\lambda^2 \exp[(2-y)2\Psi] \bar{N}^2 F + (\dot{\Psi})^2 + \epsilon \dot{\Psi}}}{\sqrt{\lambda^2 \exp[(2-y)2\Psi] \bar{N}^2 F + (\dot{\Psi})^2 - \epsilon \dot{\Psi}}} \right| \right], \end{aligned} \quad (4.36)$$

possesses the variational principle we desire. In the reduced action $\bar{N}$ is positive and independent, and $F = 1 - 2(\alpha \lambda)^{-1} e^{y\Psi} \mathbf{m}$. Also, in the expression for $I_\dagger^I$ the t' and t'' represent integration parameters rather than spacelike slices as they did before.

V. THE THERMODYNAMICAL ACTION

For both the Schwarzschild and the Witten black hole we are interested in applying the canonical action principle to a static exterior region with spatial boundary (*including* the bifurcation point) of the relevant Kruskal diagram. Such an application of the action principle is the appropriate one for studying the equilibrium thermodynamics of black holes. [5] In such a scenario, with the covariant form of the action there is no inner boundary, since the bifurcation point is a set of measure zero in the integration over $\mathcal{M}$. Nevertheless, in the canonical picture the bifurcation point is a boundary point of every spacelike slice, which implies that the canonical coordinates must obey certain fall-off conditions as the

bifurcation point is approached. Moreover, in the thermodynamical paradigm, when the initial and final spacelike slice are identified, one must worry about *regularity conditions* at the bifurcation point which ensure that the geometry is smooth. [12,5,20] To handle these issues, we will use a technique due to Brown and York. [21] The basic idea is to work with an inner boundary, but with boundary conditions which effectively seal it. The main ingredient in this technique is a new action functional, which differs from (3.1) by boundary terms. The purpose of this section is to introduce this new action principle and to study its canonical reduction via the new canonical variables.

A. Alternative canonical action principle

Starting with the canonical action (4.5), we define the new action

$$\mathcal{S}_*^I \equiv \mathcal{S}^I - \frac{1}{2} \int_{\bar{\mathcal{T}}_i} dt \bar{N} \bar{S} + \frac{1}{2} \alpha e^{-2\Phi} \eta \Big|_{B'_i}^{B''_i}. \quad (5.1)$$

Note that only inner boundary and inner corner terms have been added to the original action. It is easy to show that the canonical form of $\mathcal{S}_*^I$ has the boundary terms

$$(\mathcal{S}_*^I)_{\partial \mathcal{M}} = \int_{\bar{\mathcal{T}}_o} dt \left[\alpha e^{-2\Phi} \eta \dot{\Phi} - \bar{N} \bar{E} \right] + \int_{\bar{\mathcal{T}}_i} dt \left[\frac{1}{2} \alpha e^{-2\Phi} \dot{\eta} - \bar{N} \left(\bar{E} + \frac{1}{2} \bar{S} \right) \right]. \quad (5.2)$$

Using the expressions (4.8a) and (4.8c), one finds that the inner boundary term is

$$(\mathcal{S}_*^I)_{\bar{\mathcal{T}}_i} = - \int_{\bar{\mathcal{T}}_i} dt N \left[E - vJ + \frac{1}{2} (S - vT) \right]. \quad (5.3)$$

It is also relatively straightforward to compute the variation of $\mathcal{S}_*^I$. With the result (4.7) it follows that the $\bar{\mathcal{T}}$ boundary and corner contributions to the variation of $\mathcal{S}_*^I$ are

$$\begin{aligned} (\delta \mathcal{S}_*^I)_{\bar{\mathcal{T}}, B'', B'} = & \\ & - \int_{\bar{\mathcal{T}}_o} dt \left[\bar{E} \delta \bar{N} + \bar{N} \bar{S} \delta \Phi - \left(\bar{N} \bar{J} + \alpha e^{-2\Phi} \dot{\Phi} \right) \delta \eta \right] + \alpha e^{-2\Phi} \eta \delta \Phi \Big|_{B'_o}^{B''_o} \\ & - \int_{\bar{\mathcal{T}}_i} dt \left[\bar{E} \delta \bar{N} + \frac{1}{2} e^{-2\Phi} \delta \left(e^{2\Phi} \bar{N} \bar{S} \right) - \left(\bar{N} \bar{J} + \alpha e^{-2\Phi} \dot{\Phi} \right) \delta \eta \right] + \frac{1}{2} \alpha e^{-2\Phi} \delta \eta \Big|_{B'_i}^{B''_i}. \end{aligned} \quad (5.4)$$

We describe the boundary conditions at the inner boundary as *completely open* because $\bar{N}$ and $e^{2\Phi} \bar{S}$ are held fixed, as opposed to closed or *microcanonical* boundary conditions characterized by fixation of the energy $\bar{E}$ and fixation of Φ (effectively the B surface area). [21] With some work and the formulae in (4.8), one can show that the inner-boundary and inner-corner-point contributions in the above variation may be combined into the expression

$$(\delta \mathcal{S}_*^I)_{\bar{\mathcal{T}}_i, B''_i, B'_i} = - \int_{\bar{\mathcal{T}}_i} dt \left[(E - vJ) \delta N + \frac{1}{2} e^{-2\Phi} \delta \left(N e^{2\Phi} S - v N e^{2\Phi} T \right) - (N/\gamma^2) J \delta \eta \right]. \quad (5.5)$$

Now we introduce fall-off conditions on the fields which seal the inner boundary. What we have in mind is a general foliation of our spatially bounded static region $\mathcal{M}$. All of the spatial slices meet at the bifurcation point, but otherwise are essentially arbitrary. Our

phase space is the set of fields $(\Lambda, P_\Lambda; \Phi, P_\Phi)$ with the appropriate fall-off conditions near the bifurcation point. The needed fall-off conditions have already been given in LW for the specific case of the Schwarzschild geometry. For convenience and without loss of generality, take the inner boundary, the bifurcation point, to be located at $r = r_i = 0$ and the outer boundary to be located at $r = r_o = 1$. The boundary conditions given in LW are the following:

$$\Lambda(t, r) = \Lambda_0(t) + O(r^2) \quad (5.6a)$$

$$\Phi(t, r) = \Phi_0(t) + \Phi_2(t)r^2 + O(r^4) \quad (5.6b)$$

$$P_\Lambda(t, r) = O(r^3) \quad (5.6c)$$

$$P_\Phi(t, r) = O(r) \quad (5.6d)$$

$$N(t, r) = N_1(t)r + O(r^3) \quad (5.6e)$$

$$N^r(t, r) = N_1^r(t)r + O(r^3), \quad (5.6f)$$

where $O(r^n)$ stands for a term whose magnitude as $r \rightarrow 0$ is bounded by r^n times a constant. Also, as $r \rightarrow 0$, the k 'th derivative of such a term is similarly bounded by r^{n-k} times a constant for $1 \leq k \leq n$. Note that the time development at the inner boundary has been arrested as the lapse vanishes there. One can show that these boundary conditions are consistent with the equations of motion. In other words, the Hamiltonian evolution preserves the above boundary conditions, provided that the initial data obeys both the above set of fall-off conditions and the constraints (4.6) on the initial spacelike slice Σ and provided that the lapse and shift also obey the above fall-off conditions.⁸ Moreover, the dynamical equation for Φ and the above fall-off conditions imply that $\dot{\Phi}_0 = 0$. Imposition of the Hamiltonian constraint (4.6a) as $r \rightarrow 0$ yields the relation

$$(\Lambda_0)^2 = -4y^{-1}\lambda^{-2}\Phi_2 \exp[(y-2)2\Phi_0], \quad (5.7)$$

which shows that Φ_2 is negative for classical solutions. Let us quickly compare this result with the Schwarzschild result found in LW. Setting $R(t, r) = R_0(t) + R_2(t)r^2 + O(r^4)$ near the bifurcation point, one finds that $\lambda R_0 = e^{-\Phi_0}$ and $\lambda R_2 = -\Phi_2 e^{-\Phi_0}$. Therefore, for the SSGR case the above expression is $(\Lambda_0)^2 = 4R_0R_2$, which is the LW result.

Application of the fall-off conditions (5.6) to the inner boundary term (5.3) shows that now the new action (5.1) has the following form:

$$\begin{aligned} \mathcal{S}_*^I = & \int_{\mathcal{M}} d^2x \left(P_\Lambda \dot{\Lambda} + P_\Phi \dot{\Phi} - N\mathcal{H} - N^r \mathcal{H}_r \right) \\ & + \int_{\bar{\mathcal{T}}_o} dt \left(\alpha e^{-2\Phi} \eta \dot{\Phi} - \bar{N} \bar{E} \right) + \frac{1}{2} \int_{t'}^{t''} dt \alpha e^{-2\Phi} (N'/\Lambda) \Big|_{r=0}, \end{aligned} \quad (5.8)$$

where in the last integral t' and t'' now represent integration parameters and not manifolds as they have before. We refer to (5.8) as the *thermodynamical action* because of its importance

⁸Note that one must appeal to (5.7) when showing that the boundary conditions (5.6) are consistent with the $\dot{P}_\Lambda$ equation of motion.

in the path-integral formulation of gravitational thermodynamics. This is the appropriate action with which to study the canonical ensemble for spherically symmetric black holes. [5] Using (5.5) in tandem with the fall-off conditions (5.6), one finds the following boundary terms in the variation of the thermodynamical action:

$$\begin{aligned}
(\delta\mathcal{S}_*^I)_{\bar{T},B'',B'} &= \frac{1}{2} \int_{t'}^{t''} dt \, \alpha e^{-2\Phi} \delta(N'/\Lambda) \Big|_{r=0} \\
&\quad - \int_{\bar{\mathcal{I}}_0} dt \left[\bar{E} \delta \bar{N} + \bar{N} \bar{S} \delta \Phi - \left(\bar{N} \bar{J} + \alpha e^{-2\Phi} \dot{\Phi} \right) \delta \eta \right] + \alpha e^{-2\Phi} \eta \delta \Phi \Big|_{B'_0}^{B''_0} .
\end{aligned} \tag{5.9}$$

For the $y = 1$ case this is precisely the action and variational principle considered in LW. Notice that we have the same boundary conditions at the outer boundary as before: $\bar{N}$ and Φ are held fixed on this surface in the variational principle associated with (5.8). As spelled out in LW, the quantity N'/Λ which is fixed at the bifurcation point in the variational principle has a direct physical interpretation. In fact, N'/Λ is the time rate of change of a certain boost parameter. Each Σ slice defines its own timelike normal u^a at the bifurcation point. As the Σ slicing develops in time this vector is continuously boosted at the rate $(N'/\Lambda)|_{r=0} = N_1/\Lambda_0$.

B. Canonical reduction of the thermodynamical action

We want a new-variable version of the thermodynamical action which is amenable to canonical reduction. From the work in §4 we already know how to modify/handle the outer boundary term when passing to the new canonical variables. Except for a minor difference, we will handle the outer boundary term in the thermodynamical action just like in the previous section. Therefore, the last section has already addressed several of the delicate issues concerning the canonical reduction of the thermodynamical action. However, there is a new feature of the thermodynamical action which we need to worry about. This new feature concerns the quantity N'/Λ which is fixed at the bifurcation point. We have already discussed why fixation of $\bar{N}$ and Φ (now only at the outer boundary) are important features of the variational principle. The boundary integral at the bifurcation point is also of importance. Indeed, in the thermodynamical paradigm black-hole entropy arises from this term. [5,20] For the Schwarzschild case LW has shown that for applications to gravitational thermodynamics it is crucial to retain fixation of N'/Λ at the bifurcation point when passing to a new-variable version of the action. We also regard this feature of the action principle as the feature of central importance which needs to be preserved. Though essentially reviewing the work of LW, this subsection shows how the Louko-Whiting formalism extends to the 2dDG case. Therefore, our discussion has relevance for the thermodynamics of pure dilaton gravity.

Let us present the quantities with which we will construct a new-variable version of the thermodynamical action. The new canonical variables are the same as before, since the transformation (4.11) remains canonical with the boundary conditions adopted for the thermodynamical action. As before, the term $-\left(\alpha e^{-2\Phi} \delta \Phi \varphi\right)'$ in the identity (4.15) gives two boundary terms. The one at the outer boundary vanishes as Φ is held fixed on this boundary

element. Moreover, φ vanishes as $r \rightarrow 0$ so the inner boundary term also vanishes. Hence, upon integration (4.15) still shows that the difference between the old and new Liouville forms is an exact form. The new shift N^Ψ is again defined by (4.25b). However, following Louko and Whiting, we define a *different* new lapse

$$\mathbf{N} = -N^M \left(\frac{yM}{\alpha} \right) \left(\frac{\alpha\lambda}{2M} \right)^{2/y}. \quad (5.10)$$

Recall that M has units of length for the SSGR case and units of inverse length for 2dDG. It is easy to see that N^M is dimensionless for both cases, and, therefore, for both cases $\mathbf{N}$ has units of inverse length. It turns out that this choice for the lapse must be made in order to ensure that we retain fixation of the boost rate N'/Λ at the bifurcation point as a feature of our variational principle. In terms of $\mathbf{N}$ the boundary lapse is given by

$$\bar{N}^2 = \left(\frac{\alpha}{yM} \right)^2 \left(\frac{2M}{\alpha\lambda} \right)^{4/y} F \mathbf{N}^2 - \lambda^{-2} \exp[(y-2)2\Psi] F^{-1} (N^\Psi)^2. \quad (5.11)$$

The fall-off conditions (5.6) imply the following fall-off conditions for the new variables:

$$M(t, r) = \frac{1}{2}\alpha\lambda \exp[-y\Psi_0(t)] + M_2(t)r^2 + O(r^4) \quad (5.12a)$$

$$\Psi(t, r) = \Psi_0(t) + \Psi_2 r^2 + O(r^4) \quad (5.12b)$$

$$P_M(t, r) = O(r) \quad (5.12c)$$

$$P_\Psi(t, r) = O(r) \quad (5.12d)$$

$$\mathbf{N}(t, r) = \mathbf{N}_0(t) + O(r^2) \quad (5.12e)$$

$$N^\Psi(t, r) = N_2^\Psi(t)r^2 + O(r^4), \quad (5.12f)$$

where

$$\Psi_0 = \Phi_0 \quad (5.13a)$$

$$\Psi_2 = \Phi_2 \quad (5.13b)$$

$$M_2 = -\frac{1}{2}\alpha\lambda\Phi_2 \exp(-y\Phi_0) \left[y + 4\lambda^{-2}(\Lambda_0)^{-2}\Phi_2 \exp[(y-2)2\Phi_0] \right] \quad (5.13c)$$

$$\mathbf{N}_0 = -\frac{1}{4}y\lambda^2 N_1 \Lambda_0 (\Phi_2)^{-1} \exp[(2-y)2\Phi_0] \quad (5.13d)$$

$$N_2^\Psi = 2\Phi_2 N_1^r. \quad (5.13e)$$

We also have that

$$F = 4\lambda^{-2}(\Phi_2)^2(\Lambda_0)^{-2} \exp[(y-2)2\Phi_0] r^2 + O(r^4). \quad (5.13)$$

For the SSGR case these fall-off results for the new variables match those given in LW. Equation (5.7) implies that

$$(N'/\Lambda)|_{r=0} = -\frac{1}{4}y\lambda^2 N_1 \Lambda_0 (\Phi_2)^{-1} \exp[(2-y)2\Phi_0], \quad (5.14)$$

which in turn gives $\mathbf{N}_0 = N_1/\Lambda_0$. Hence, we want to ensure that the Lagrange multiplier $\mathbf{N}$ is fixed at the bifurcation point in our new variational principle.

Let us now consider the new-variable version of the thermodynamical action and show that it has the correct variational principle. We could write down a general action which covers both the SSGR and 2dDG cases, but the expression is a bit unseemly. Therefore, let us examine both cases separately. For the SSGR case we have

$$\begin{aligned} \mathcal{S}_\diamond^I \equiv & \int_{\mathcal{M}} d^2x \left(P_M \dot{M} + P_R \dot{R} + 4NMM' - N^R P_R \right) \\ & + \int_{t'}^{t''} dt \left. 2M^2 \mathbf{N} \right|_{r=0} + \int_{\bar{\mathcal{T}}_o} dt \left[-\rho R \dot{R} - \bar{N} \bar{E} \right], \end{aligned} \quad (5.15)$$

with the boundary lapse given by

$$\bar{N}^2 = 16M^2 F \mathbf{N}^2 - F^{-1} \left(N^R \right)^2. \quad (5.16)$$

To get these expressions we have used the facts that $e^{-\Psi} = \lambda R$, $P_\Psi = -R P_R$, and $N^\Psi = -N^R/R$ (all in the notation of KVK and LW). Also, $\bar{E}$ and ρ are still given by (4.31b) and (4.32), respectively, but now one must express them in terms of R . This is precisely the action considered in LW. For Witten's 2dDG model we find

$$\begin{aligned} \mathcal{S}_\diamond^I \equiv & \int_{\mathcal{M}} d^2x \left[P_M \dot{M} + P_\Psi \dot{\Psi} + \lambda^{-1} N M' - N^\Psi P_\Psi \right] \\ & + \int_{t'}^{t''} dt \left. \lambda^{-1} M \mathbf{N} \right|_{r=0} + \int_{\bar{\mathcal{T}}_o} dt \left[\alpha e^{-2\Psi} \rho \dot{\Psi} - \bar{N} \bar{E} \right], \end{aligned} \quad (5.17)$$

with

$$\bar{N}^2 = \lambda^{-2} F \mathbf{N}^2 - \lambda^{-2} F^{-1} \left(N^\Psi \right)^2. \quad (5.18)$$

In contrast to the SSGR case, for 2dDG the Lagrange multiplier $N^M|_{r=0}$ does, apart only from a dimensionful constant, specify the boost rate of the Σ normal u^a at the bifurcation point.

The variation of (5.15) has already been considered in LW, so we will only consider the variation of (5.17). The equations of motion derived from (5.17) are the same as those given in (4.33), except that now $\dot{P}_M = -\lambda^{-1} \mathbf{N}$. Upon variation of the action (5.17), we find the boundary terms

$$\begin{aligned} (\delta \mathcal{S}_\diamond^I)_{\partial \mathcal{M}} = & \int_{t'}^{t''} dr \left(P_M \delta M + P_\Psi \delta \Psi \right) + \int_{t'}^{t''} dt \left. \lambda^{-1} M \delta \mathbf{N} \right|_{r=0} \\ & + \int_{\bar{\mathcal{T}}_o} dt \left(\bar{\Pi}_N \delta \bar{N} + \bar{\Pi}_\Psi \delta \Psi + \bar{\Pi}_M \delta M \right) + \alpha e^{-2\Psi} \rho \delta \Psi \Big|_{B'_o}^{B''_o}. \end{aligned} \quad (5.19)$$

In the first integral t' and t'' represent spacelike slices, while in the second integral they are integration parameters. The $\bar{\mathcal{T}}_o$ momenta in (5.19) are essentially the same as the outer-boundary ones in (4.35), except that now $\bar{N}$ stands for (5.18) and the momenta $\bar{\Pi}_M = \lambda^{-1} \mathbf{N} + (\alpha \lambda)^{-1} e^{y\Psi} F^{-1} \bar{N} \bar{E}$. Like before, plugging in the explicit formula (5.18) for $\bar{N}^2$, one finds that $\bar{\Pi}_M$ vanishes when the equation of motion $\dot{P}_M = -\lambda^{-1} \mathbf{N}$ holds. Therefore, M need not be held fixed on $\bar{\mathcal{T}}_o$ in the variational principle associated with $\mathcal{S}_\diamond^I$. It is now straightforward to pass to the reduced form of the thermodynamical action.

VI. DISCUSSION

We conclude with some comments concerning the possible extension of the KVK and LW formalisms to other two-dimensional models of gravity. Recently, important progress has been made in the field of two-dimensional gravity with the realization that a huge class of two-dimensional models can be described within the framework of the so-called *Poisson-sigma models* (PSM's) of Klösch, Schaller, and Strobl. [22] For all such models there exists an absolutely conserved quantity C (referred to as a *Casimir function* in the Poisson-sigma model language) which is analogous to our M expression (3.16), and recently Kummer and Widerin have explored the relationship between the PSM C and notions of quasilocal energy for such models. [23] Many of our results, especially those concerning our general treatment of quasilocal energy-momentum, seem to extend to the general PSM formalism. In particular, the absolutely conserved quantity C can be interpreted as a quasilocal boost invariant. [24] Extension of the canonical-reduction method of KVK to PSM theory also seems possible, though several technical difficulties lie in the way. For instance, one encounters an almost limitless variety of singularity structures when considering the set of all PSM's. [22] For SSGR the canonical transformation of KVK is singular at the horizon. Similar technical difficulties are likely to surface when applying the KVK method to any two-dimensional model. Since the collection of all possible Penrose diagrams obtainable from PSM's is so large, it is questionable whether or not a fully unified treatment for the canonical reduction of all PSMs is possible. On the other hand, the richness of singularity structures in PSM gravitation offers what is perhaps a promising testing ground for gravitational thermodynamics. The appropriate thermodynamical action, as expressed in the LW formalism, would be a crucial ingredient in any study of PSM thermodynamics via reduced canonical variables.

VII. ACKNOWLEDGMENTS

For discussions and helpful comments I thank H. Balasin, J. D. Brown, W. Kummer, F. Schramm, S. Sinha, and J. W. York. The bulk of this research has been supported by the Fonds zur Förderung der wissenschaftlichen Forschung (Lise Meitner Fellowship M-00182-PHY). Also, some of this work was done while visiting the University of North Carolina at Chapel Hill with support provided by the National Science Foundation, grant number PHY-9413207. Finally, this work was begun in Pune at the Inter-University Centre for Astronomy & Astrophysics, with support from the Indo-US Exchange Programme and the University Grants Commission of India.

NOTE ADDED: After this project was completed, we learned of a new paper [25] by Varadarajan which also treats the canonical reduction of two-dimensional pure dilaton gravity in the manner of KVK. Moreover, this paper also considers quantization.

APPENDIX A: REDUCTION (BY SPHERICAL SYMMETRY) OF THE HILBERT ACTION WITH BOUNDARY TERMS

Take our two-dimensional metric g_{ab} from the preliminary section and adjoin to it the metric of a round sphere. The result is the four-dimensional spherically symmetric metric

$${}^4g_{\mu\nu}dx^\mu dx^\nu = -N^2 dt^2 + \Lambda^2 (dr + N^r dt)^2 + R^2 (d\theta^2 + \sin^2 \theta d\phi^2). \quad (\text{A1})$$

Now every point $B(r, t)$ of $\mathcal{M}$ is a round sphere of radius $R(t, r)$. Therefore, our $1 + 1$ spacetime region $\mathcal{M}$ has now been promoted to a four-dimensional time history ${}^4\mathcal{M}$ of a three-dimensional spatial region which lies between concentric spheres. Slices Σ and sheets $\bar{\mathcal{T}}$ now correspond to three-dimensional hypersurfaces of ${}^4\mathcal{M}$. In particular, the boundary elements $\bar{\mathcal{T}}_i$ and $\bar{\mathcal{T}}_o$ are now $2 + 1$ hypersurfaces in ${}^4\mathcal{M}$.

The action functional associated with ${}^4\mathcal{M}$ is the standard Einstein-Hilbert action, complete with the TrK terms needed to ensure that the four-dimensional action principle features fixation of the induced three-metric on all of the elements of the three-boundary $\partial^4\mathcal{M}$. [14,10] In this appendix we will insert the metric ansatz (A1) into the four-dimensional spacetime action and then integrate out the angular variables. This procedure yields a “spherically reduced” action principle for the fields g_{ab} and R (or equivalently Φ) which are defined on the toy $1 + 1$ dimensional spacetime described in the preliminary section. The variational principle for the spherically reduced action still features fixation of the induced metric on the boundary, and the equations of motion derived from the reduced action are the four-dimensional Einstein equations subject to the ansatz of spherical symmetry; however, we do not prove this latter claim here. Ref. [13] noted for the first time that the spherically reduced action reproduces the correct four-dimensional Einstein equations.

The four-dimensional spacetime action is [14,10]

$$16\pi\mathcal{S}^I = \int_{{}^4\mathcal{M}} d^4x \sqrt{-{}^4g} \mathfrak{R} + 2 \int_{t'}^{t''} d^3x \sqrt{h} K - 2 \int_{\bar{\mathcal{T}}} d^3x \sqrt{-\bar{\gamma}} \bar{\Theta} - 2 \int_{B'}^{B''} d^2x \sqrt{\sigma} \eta, \quad (\text{A2})$$

where $\mathfrak{R}$ is the Ricci scalar of ${}^4\mathcal{M}$, h_{ij} is the induced three-metric on the spacelike slices t' and t'' , and $\bar{\gamma}_{ij}$ is the induced three-metric on the $\bar{\mathcal{T}}$ boundary elements. In (A2) we are using integral notations similar to those listed in (3.2). Also, $K_{\mu\nu} \equiv -h_\mu^\lambda {}^4\nabla_\lambda u_\nu$ is the extrinsic curvature of t' or t'' as embedded in ${}^4\mathcal{M}$, where ${}^4\nabla_\mu$ is the torsion-free covariant derivative operator compatible with ${}^4g_{\mu\nu}$. In this definition u_μ is the future-pointing normal of t' or t'' , and we have used spacetime coordinates for convenience, so $h_{\mu\nu} = {}^4g_{\mu\nu} + u_\mu u_\nu$. Likewise, the extrinsic curvature of the $\bar{\mathcal{T}}$ boundary as embedded in ${}^4\mathcal{M}$ is $\bar{\Theta}_{\mu\nu} \equiv -\bar{\gamma}_\mu^\lambda {}^4\nabla_\lambda \bar{n}_\nu$, where $\bar{n}_\mu$ is the outward-pointing spacelike normal of $\bar{\mathcal{T}}$ boundary and here $\bar{\gamma}_{\mu\nu} = {}^4g_{\mu\nu} - \bar{n}_\mu \bar{n}_\nu$. Finally, on the corners B' and B'' (each is the disjoint union of an inner and outer sphere) the two-metric is σ_{ab} and $\eta = \sinh^{-1}(u_\mu \bar{n}^\mu)$.

Let us collect a few results needed for the reduction by spherical symmetry. The letter Σ denotes the constant time slices associated with the line-element (A1), and the set of nonzero components of the Σ extrinsic curvature tensor K_j^i is the following:

$$K_r^r = -(N\Lambda)^{-1} [\dot{\Lambda} - (\Lambda N^r)'] \quad (\text{A3a})$$

$$K_\theta^\theta = K_\phi^\phi = -(NR)^{-1} (\dot{R} - N^r R') \quad (\text{A3b})$$

Treating the time-radial piece g_{ab} of the full four-metric (A1) as if it were a true metric in its own right, we can compute its curvature scalar $\mathcal{R}[g]$. The result is

$$\mathcal{R} = -2(N\Lambda)^{-1} (\Lambda K_r^r)^\bullet - 2(N\Lambda)^{-1} (\Lambda^{-1} N' - \Lambda N^r K_r^r)'. \quad (\text{A4})$$

Now we turn to the reduction. Let us start with the spacetime volume integral in (A2). Consider the identity [14]

$$\mathfrak{R}[^4g] = {}^3R[h] + K_{\mu\nu}K^{\mu\nu} - (K)^2 - 2 {}^4\nabla_\mu (Ku^\mu + a^\mu) , \quad (\text{A5})$$

where a^μ is the spacetime acceleration of u^μ and 3R stands for the Σ curvature scalar (not to be confused with the radius function R). Use this identity to split the volume integral in (A2) into three pieces:

$$(\text{term 1}) = \int_{^4\mathcal{M}} d^4x \sqrt{-^4g} {}^3R \quad (\text{A6a})$$

$$(\text{term 2}) = \int_{^4\mathcal{M}} d^4x \sqrt{-^4g} [K_{ij}K^{ij} - (K)^2] \quad (\text{A6b})$$

$$(\text{term 3}) = -2 \int_{^4\mathcal{M}} d^4x \partial_\mu [\sqrt{-^4g} (Ku^\mu + a^\mu)] . \quad (\text{A6c})$$

Focus attention on

$$(\text{term 3}) = -2 \int_{^4\mathcal{M}} d^4x \left\{ [\sqrt{-^4g} (Ku^t + a^t)]^\bullet + [\sqrt{-^4g} (Ku^r + a^r)]' \right\} , \quad (\text{A7})$$

where one has

$$u^t = 1/N \quad (\text{A8a})$$

$$u^r = -N^r/N \quad (\text{A8b})$$

$$a^t = 0 \quad (\text{A8c})$$

$$a^r = N^{-1}\Lambda^{-2}N' . \quad (\text{A8d})$$

After some work and with $\sqrt{-^4g} = N\Lambda R^2 \sin \theta$, this term can be written as

$$\begin{aligned} (\text{term 3}) = & \int_{^4\mathcal{M}} d^4x \sqrt{-^4g} [\mathcal{R} + 4K_r^r K_\theta^\theta - 4\Lambda^{-2}(\log N)'(\log R)'] \\ & + 4 \int_{^4\mathcal{M}} d^4x \sin \theta \left[(\Lambda R^2 N^r K_\theta^\theta)' - (\Lambda R^2 K_\theta^\theta)^\bullet \right] , \end{aligned} \quad (\text{A9})$$

where we have used $K_\theta^\theta = K_\phi^\phi$. Straightforward manipulations show that

$$(\text{term 2}) = \int_{^4\mathcal{M}} d^4x \sin \theta [-2K_\theta^\theta K_\theta^\theta - 4K_r^r K_\theta^\theta] . \quad (\text{A10})$$

Recombining the three terms, we get

$$\begin{aligned} \int_{^4\mathcal{M}} d^4x \sqrt{-^4g} \mathfrak{R} = & \int_{^4\mathcal{M}} d^4x \sqrt{-^4g} [\mathcal{R} + {}^3R - 2K_\theta^\theta K_\theta^\theta - 4\Lambda^{-2}(\log N)'(\log R)'] \\ & + 4 \int_{^4\mathcal{M}} d^4x \sin \theta \left[(\Lambda R^2 N^r K_\theta^\theta)' - (\Lambda R^2 K_\theta^\theta)^\bullet \right] . \end{aligned} \quad (\text{A11})$$

Now the explicit expression for the Σ Ricci scalar is

$${}^3R = -4\Lambda^{-2}R^{-1}R'' + 4\Lambda^{-3}R^{-1}\Lambda'R' - 2\Lambda^{-2}R^{-2}(R')^2 + 2R^{-2} . \quad (\text{A12})$$

With this expression and the result for K_θ^θ , one can show that

$${}^3R - 2K_\theta^\theta K_\theta^\theta = 2R^{-2} \left[g^{ab} \nabla_a R \nabla_b R + 1 - 2\Lambda^{-1} (\Lambda^{-1} R R')' \right]. \quad (\text{A13})$$

Therefore, after a bit of re-shuffling, one finds the following expression for the spacetime volume term in the action:

$$\begin{aligned} \int_{4\mathcal{M}} d^4x \sqrt{-4g} \mathfrak{R} &= \int_{4\mathcal{M}} d^4x \sqrt{-4g} \left[\mathcal{R} + 2R^{-2} g^{ab} \nabla_a R \nabla_b R + 2R^{-2} \right] \\ &\quad + 4 \int_{4\mathcal{M}} d^4x \sin \theta \left[(\Lambda R^2 N^r K_\theta^\theta - \Lambda^{-1} N R R')' - (\Lambda R^2 K_\theta^\theta)^\bullet \right]. \end{aligned} \quad (\text{A14})$$

Now we turn to the boundary terms in the action. First consider

$$2 \int_{t'}^{t''} d^3x \sqrt{h} K = 2 \int_{t'}^{t''} d^3x \sqrt{h} K_r^r + 4 \int_{t'}^{t''} d^3x \Lambda R^2 \sin \theta K_\theta^\theta. \quad (\text{A15})$$

For the $\bar{\mathcal{T}}$ boundary terms we have

$$-2 \int_{\bar{\mathcal{T}}} d^3x \sqrt{-\bar{\gamma}} \bar{\Theta} = -2 \int_{\bar{\mathcal{T}}} d^3x \sqrt{-\bar{\gamma}} \bar{\Theta}_t^t - 2 \int_{\bar{\mathcal{T}}} d^3x \sqrt{-\bar{\gamma}} \sigma_{ij} \bar{\Theta}^{ij}, \quad (\text{A16})$$

where $\sigma_{ij} \bar{\Theta}^{ij} = \bar{\Theta}_\theta^\theta + \bar{\Theta}_\phi^\phi$. Now use the result⁹

$$\sigma_{ij} \bar{\Theta}^{ij} = \gamma k + v \gamma \sigma_{ij} K^{ij}. \quad (\text{A17})$$

For the case at hand, $\sigma_{ij} K^{ij} = 2K_\theta^\theta$ and

$$k = \pm 2\Lambda^{-1} R^{-1} R' \quad (\text{A18})$$

with $-$ at the outer sphere and $+$ at the inner sphere. Therefore, the $\bar{\mathcal{T}}$ boundary integrals can be expressed as

$$-2 \int_{\bar{\mathcal{T}}} d^3x \sqrt{-\bar{\gamma}} \bar{\Theta} = -2 \int_{\bar{\mathcal{T}}} d^3x \sqrt{-\bar{\gamma}} \bar{\Theta}_t^t + 4 \int_{\bar{\mathcal{T}}_i}^{\bar{\mathcal{T}}_o} d^3x N \sin \theta \left[\Lambda^{-1} R R' - \mathbf{v} R^2 K_\theta^\theta \right]. \quad (\text{A19})$$

To get (A19), we have used $\sqrt{-\bar{\gamma}} \gamma = N R^2 \sin \theta$ (note that the γ 's on the left-hand side are different, the second is the relativistic factor). Also, in the above expression

$$\int_{\bar{\mathcal{T}}_i}^{\bar{\mathcal{T}}_o} = \int_{\bar{\mathcal{T}}_o} - \int_{\bar{\mathcal{T}}_i}, \quad (\text{A20})$$

and $\mathbf{v} = \Lambda N^r / N$. The v in (A17) is $-\mathbf{v}$ on $\bar{\mathcal{T}}_i$ and $\mathbf{v}$ on $\bar{\mathcal{T}}_o$.

⁹Proving this result is an exercise with projection operators. Note that the extrinsic curvature of B as embedded in Σ is defined in spacetime coordinates by $k_{\mu\nu} = -\sigma_\mu^\lambda D_\lambda n_\nu$, where D_μ is the Σ intrinsic covariant derivative operator compatible with $h_{\mu\nu}$ and $\sigma_{\mu\nu} = {}^4g_{\mu\nu} + u_\mu u_\nu - n_\mu n_\nu$. On a Σ covector like n_ν , the action of D_μ is $D_\mu n_\nu = h_\mu^\lambda h_\nu^{\kappa 4} \nabla_\lambda n_\kappa$. Also remember that $\bar{n}^\mu = \gamma n^\mu + v \gamma u^\mu$.

Using the results (A14), (A15), and (A19), we see that with the metric ansatz (A1) the spacetime action (A2) reduces to

$$16\pi\mathcal{S}^I = \int_{\mathcal{M}} d^4x \sqrt{-g} \left[\mathcal{R} + 2R^{-2} g^{ab} \nabla_a R \nabla_b R + 2R^{-2} \right] \\ + 2 \int_{t'}^{t''} d^3x \sqrt{h} K_r^r - 2 \int_{\bar{T}} d^3x \sqrt{-\bar{\gamma}} \bar{\Theta}_t^t - 2 \int_{B'}^{B''} d^2x \sqrt{\sigma} \eta. \quad (\text{A21})$$

Now define $\mathcal{K} \equiv K_r^r$ and $\bar{\vartheta} \equiv \bar{\Theta}_t^t$. None of the quantities in the above action has any angular dependence. Therefore, we simply integrate over the angular variables to find

$$\mathcal{S}^I = \frac{1}{4} \int_{\mathcal{M}} d^2x \sqrt{-g} R^2 \left[\mathcal{R} + 2R^{-2} g^{ab} \nabla_a R \nabla_b R + 2R^{-2} \right] \\ + \frac{1}{2} \int_{t'}^{t''} dr \Lambda R^2 \mathcal{K} - \frac{1}{2} \int_{\bar{T}} dt \bar{N} R^2 \bar{\vartheta} - \frac{1}{2} R^2 \eta \Big|_{B'}^{B''}. \quad (\text{A22})$$

If one prefers, one can write the action in terms of the dilaton $\Phi = -\log \lambda R$. The result is precisely the action (3.1) from the first section with the choices $y = 1$ and $\alpha = \lambda^{-2}$.

APPENDIX B: THE WITTEN BLACK HOLE

In this appendix we have two goals in mind. The first is to compute the energy at spatial infinity (associated with the static-time slices $\tilde{\Sigma}$) for the Witten black hole. The second is to derive expressions, depending on the canonical variables of an arbitrary spacelike slice Σ , for the Witten-black-hole mass parameter M_w and (minus) the radial derivative of the Killing time $-\tau'$. Our procedure for obtaining such expressions is nearly identical to one found in KVK. Multiplying the expression we find for M_w by $\alpha/2$, one finds the boost invariant M given in (3.16).

1. Line-element and asymptotic energy

In Kruskal-type coordinates the black-hole solution of (vacuum) 2dDG is given by the line-element

$$ds^2 = - \left(M_w / \lambda - \lambda^2 x^+ x^- \right)^{-1} dx^+ dx^-, \quad (\text{B1})$$

along with the following expression for the dilaton:

$$e^{-2\Phi} = M_w / \lambda - \lambda^2 x^+ x^-. \quad (\text{B2})$$

Our form of the line-element corresponds to the one given in Ref. [26], and the discussion which follows uses the conventions of that reference. We are only interested in the right static region of the Kruskal diagram associated with (B1).

We wish to compute the energy at infinity associated with the preferred static-time slices $\tilde{\Sigma}$, so we first have to find these slices. Consider the new coordinates (τ, σ) defined by [26]

$$\lambda x^- = -e^{-\lambda(\tau-\sigma)} \quad (\text{B3a})$$

$$\lambda x^+ = e^{\lambda(\tau+\sigma)} . \quad (\text{B3b})$$

Note that the coordinate patch (τ, σ) only covers the right static region of the Kruskal diagram. In terms of these coordinates the line-element reads

$$ds^2 = \left(1 + e^{-2\lambda\sigma} M_w / \lambda\right)^{-1} \left(-d\tau^2 + d\sigma^2\right) . \quad (\text{B4})$$

Notice that as $\sigma \rightarrow \infty$ the line element becomes flat. Further, notice that as $\sigma \rightarrow \infty$ the dilaton behaves as

$$\Phi = -\lambda\sigma - e^{-2\lambda\sigma} M_w / 2\lambda + O(e^{-2\lambda\sigma}) , \quad (\text{B5})$$

where $O(e^{-2\lambda\sigma})$ stands for quadratic and higher powers in $e^{-2\lambda\sigma}$. At spatial infinity the black-hole solution approaches the *linear dilaton*, the vacuum solution of the theory. For the linear dilaton $\Phi = -\lambda\sigma$ and the line element is Minkowskian.

The static-time slices $\tilde{\Sigma}$ are level surfaces of constant τ . We shall pick one and evaluate the energy at an outer boundary point B_o . Now the normal of such a point as embedded in $\tilde{\Sigma}$ has been denoted $\tilde{\mathbf{n}}^a$. Since only the static-time slices are of interest now, we shall drop the tilde which appears on $\tilde{\Sigma}$, $\tilde{\mathbf{n}}^a$, and other objects. Furthermore, let us simply write n^a for $\mathbf{n}^a$, with the understanding that the rest of our analysis in this appendix deals exclusively with an *outer* boundary point. The history of B_o with respect to the Eulerian observers of the Killing time slices is the timelike sheet $\mathcal{T}_o$ defined by $\sigma = \sigma_o \geq -\infty$, where σ_o is a finite constant. This means that the dilaton Φ is also fixed to a constant value Φ_o on $\mathcal{T}_o$. Now it turns out that the energy expression for B_o as embedded in the static slice diverges in the limit that $\sigma_o \rightarrow \infty$.

In order to obtain a finite energy at spatial infinity, we must reference the energy against the linear dilaton vacuum *before* taking the limit. The expression for the quasilocal energy *with reference point* is

$$E = \alpha e^{-2\Phi} \left[n^a \nabla_a \Phi - (n^a \nabla_a \Phi)^0 \right] . \quad (\text{B6})$$

This expression has been obtain by comparison with the known expression for quasilocal energy in general relativity, [6]

$$E = (8\pi)^{-1} \int_B d^2x \sqrt{\sigma} (k - k^\theta) . \quad (\text{B7})$$

In this expression $\sqrt{\sigma}$ is the square root of the determinant of the B metric (in our case that of a round sphere, $R^2 \sin \theta$), k is the trace of the extrinsic curvature of B as embedded in the three-dimensional hypersurface Σ of interest, and k^θ is the trace of the extrinsic curvature of a two-surface isometric to B which is embedded in three-dimensional flat Euclidean space.¹⁰

¹⁰Such a construction is not possible for a generic two surface. Of course, such a construction is always possible when B is a round sphere, the relevant case for this work.

The origin of the reference term k^0 can be traced to the freedom to add a subtraction term (a functional of the fixed boundary data) to the four-dimensional spacetime action without affecting the variational principle. [6] Likewise, the reference point contribution in (B6) arises from the freedom to append a subtraction term $-\mathcal{S}^0$ to our base action (3.1). By inspecting (B7), we see that the correct way to calculate the referenced energy is to first calculate $n^a \nabla_a \Phi - (n^a \nabla_a \Phi)|^0$ and then multiply by the nonlinear “determinant” factor $\alpha e^{-2\Phi}$.

We shall compute the quasilocal energy for the black hole with the subtraction term $(n^a \nabla_a \Phi)|^0$ determined by the linear-dilaton vacuum. For the black-hole solution the outward-pointing normal to points embedded in the constant τ slices is

$$n^a \partial / \partial x^a = \left(1 + e^{-2\lambda\sigma} M_w / \lambda\right)^{1/2} \partial / \partial \sigma. \quad (\text{B8})$$

With this one finds

$$n^a \nabla_a \Phi = -\lambda \left(1 + e^{-2\lambda\sigma} M_w / \lambda\right)^{-1/2}. \quad (\text{B9})$$

A similar calculation for the case of the linear dilaton gives

$$n^a \nabla_a \Phi|^0 = -\lambda. \quad (\text{B10})$$

Hence, for the point B_o located at a $\sigma = \sigma_o$ as embedded in the constant- τ slice, the associated referenced quasilocal energy is

$$E = \alpha \lambda \left[e^{2\lambda\sigma} + M_w / \lambda \right] \left[1 - \left(1 + e^{-2\lambda\sigma} M_w / \lambda\right)^{-1/2} \right] \Big|_{\sigma=\sigma_o}. \quad (\text{B11})$$

We then have that $\lim_{\sigma_o \rightarrow \infty} E = \frac{1}{2} \alpha M_w$. Hence we obtain the on-shell value of M given in (3.16) as the asymptotic energy associated with the static time slices. Note that the asymptotic energy is M_w if we make the choice $\alpha = 2$ for 2dDG.

2. Canonical expressions for M_w and $-\tau'$

Use of the dilaton itself as the radial coordinate casts the line-element (B4) in the Schwarzschild-like form

$$ds^2 = -F d\tau^2 + F^{-1} (d\Phi / \lambda)^2, \quad (\text{B12})$$

where $F \equiv 1 - e^{2\Phi} M_w / \lambda$. The horizon is located at $\Phi = -(1/2) \log(M_w / \lambda)$, or equivalently at $R = \sqrt{M_w / \lambda^3}$, and we know from the Kruskal form of the line-element (B1) that the geometry is perfectly regular at the horizon ($x^+ x^- = 0$). The goal now is to obtain canonical expressions for τ' and M_w .

To get the desired expressions follow the method of KVK and assume that $\tau = \tau(t, r)$ and $\Phi = \Phi(t, r)$. It proves convenient to define a dimensionful dilaton $\bar{\Phi} \equiv \Phi / \lambda$. Now expand the differentials $d\tau = \dot{\tau} dt + \tau' dr$ and $d\bar{\Phi} = \dot{\bar{\Phi}} dt + \bar{\Phi}' dr$ and plug these into the line-element (B12). Comparison of the result with the ADM form of the metric (2.1) gives the following equations:

$$\Lambda^2 = -F(\tau')^2 + F^{-1}(\bar{\Phi}')^2 \quad (\text{B13a})$$

$$\Lambda^2 N^r = -F\dot{\tau}\tau' + F^{-1}\dot{\bar{\Phi}}\bar{\Phi}' \quad (\text{B13b})$$

$$-N^2 + (\Lambda N^r)^2 = -F(\dot{\tau})^2 + F^{-1}(\dot{\bar{\Phi}})^2. \quad (\text{B13c})$$

From these it is straightforward to obtain the following expressions for the lapse and shift:

$$N^r = \frac{-F\dot{\tau}\tau' + F^{-1}\dot{\bar{\Phi}}\bar{\Phi}'}{-F(\tau')^2 + F^{-1}(\bar{\Phi}')^2} \quad (\text{B14a})$$

$$N = \frac{\dot{\bar{\Phi}}\tau' - \dot{\tau}\bar{\Phi}'}{\sqrt{-F(\tau')^2 + F^{-1}(\bar{\Phi}')^2}}. \quad (\text{B14b})$$

In obtaining the formula for N , we have taken a square root. Therefore, we need to verify that we have taken this root in such a way that the lapse is positive in the right static region of the Kruskal diagram, since we want our spacelike slices to advance everywhere into the future. Note that the dilaton is a “bad” radial coordinate in the sense that $\Phi \rightarrow -\infty$ as one approaches spatial infinity, whereas the preliminary section has assumed that the radial coordinate r increases in the direction of spatial infinity. Therefore, in the right static region $t = \tau$ and $r = -\bar{\Phi}$ are “good” coordinates, and, using these, we see that the lapse is positive everywhere in the right static region. The next step is to insert the last two expressions into the formula for the momentum

$$P_\Lambda = \alpha e^{-2\Phi} N^{-1} (\dot{\Phi} - N^r \Phi'). \quad (\text{B15})$$

After some algebra this insertion yields the first of our desired expressions

$$-\tau' = (\alpha\lambda)^{-1} e^{2\Phi} F^{-1} \Lambda P_\Lambda. \quad (\text{B16})$$

Now, using this expression for $-\tau'$ in the first equation of (B13), we find the canonical expression for F ,

$$F = (\alpha\lambda)^{-2} \left[(\alpha\Phi'/\Lambda)^2 - (e^{2\Phi} P_\Lambda)^2 \right]. \quad (\text{B17})$$

Solving for M_w , one finds the second desired expression,

$$M_w = (\alpha\lambda)^{-2} \lambda e^{2\Phi} (P_\Lambda)^2 - \lambda^{-1} e^{-2\Phi} (\Phi'/\Lambda)^2 + \lambda e^{-2\Phi}. \quad (\text{B18})$$

Notice that $M = \frac{1}{2}\alpha M_w$ is precisely the boost invariant (3.16) with the appropriate choices for 2dDG.

REFERENCES

- [1] K. V. Kuchař, Phys. Rev. **D50**, 3961 (1994).
- [2] R. Arnowitt, S. Deser, and C. W. Misner, in *Gravitation: An Introduction to Current Research*, edited by L. Witten (Wiley, New York, 1962).
- [3] T. Thiemann and H. A. Kastrup, Nucl. Phys. **B399**, 211 (1993); T. Thiemann and H. A. Kastrup, Nucl. Phys. **B399**, 221 (1993); H. A. Kastrup and T. Thiemann, Nucl. Phys. **B425**, 665 (1994); T. Thiemann, Nucl. Phys. **B436**, 681 (1995).
- [4] J. Louko and B. F. Whiting, Phys. Rev. **D51**, 5583 (1995).
- [5] J. W. York, Phys. Rev. **D33**, 2092 (1986); H. W. Braden, B. F. Whiting, and J. W. York, Phys. Rev. **D36**, 3614 (1987); B. F. Whiting and J. W. York, Phys. Rev. Lett. **61**, 1336 (1988); J. W. York, Physica **A158**, 425 (1989); J. D. Brown, G. L. Comer, E. A. Martinez, J. Melmed, B. F. Whiting, and J. W. York, Class. Quantum Grav. **7**, 1433 (1990); H. W. Braden, J. D. Brown, B. F. Whiting, and J. W. York, Phys. Rev. **D42**, 3376 (1990); J. W. York in *Conceptual Problems of Quantum Gravity*, edited by A. Ashtekar and J. Stachel (Birkhäuser, Boston, 1991); J. D. Brown, E. A. Martinez, and J. W. York, Phys. Rev. Lett. **66**, 2281 (1991).
- [6] J. D. Brown and J. W. York, Phys. Rev. **D47**, 1407 (1993).
- [7] J. D. Brown, S. R. Lau, and J. W. York, Jr., *General spacetime kinematics, the gravitational action, and quasilocal stress-energy-momentum*, manuscript in preparation, 1996.
- [8] E. Witten, Phys. Rev. **D44**, 314 (1991).
- [9] C.G. Callen, S. B. Giddings, J. A. Harvey, and A. Strominger, Phys. Rev. **D45**, R1005 (1992).
- [10] G. Hayward and K. Wong, Phys. Rev. **D46**, 620 (1992); G. Hayward, Phys. Rev. **D47**, 4778 (1993).
- [11] R. B. Mann, A. Shiekh and L. Tarasov, Nucl. Phys. **B341** 134 (1990); D. Banks and M. O’Loughlin, Nucl. Phys. **B362** 649 (1991); S. D. Odintsov and I. J. Shapiro, Phys. Lett. **B263** 183 (1991), Mod. Phys. Lett. **A7** 437 (1992); I. G. Russo and A. A. Tseytlin, Nucl. Phys. **B382** 259 (1992); I. V. Volovich, Mod. Phys. Lett. **A**, 1827 (1992); R. B. Mann, Phys. Rev. **D47** 4438 (1993); D. Louis-Martinez, J. Gegenberg, and G. Kunstatter, Phys. Rev. **D49** 5227 (1994); J. S. Lemos and P. M. Sa, Phys. Rev. **D49** 2897 (1994); M. O. Katanaev, W. Kummer and H. Liebl, *Geometric interpretation of generalized dilaton gravity*, TUW-95-09, Vienna, May 1995, gr-qc/9511009 (to appear in Phys. Rev. **D**).
- [12] G. W. Gibbons and S. W. Hawking, Commun. Math. Phys. **66**, 291 (1979).
- [13] P. Thomi, B. Isaak, and P. Hájíček, Phys. Rev. **D30**, 1168 (1984).
- [14] J. W. York, Phys. Rev. Lett. **28**, 1082 (1972); J. W. York, Found. Phys. **16**, 249 (1986).
- [15] S. R. Lau, *New variables, the gravitational action, and boosted quasilocal stress-energy-momentum*, IUCAA 34/94, December 1994 and TUW-95-18, July 1995, gr-qc/9504026 (to appear in Class. Quantum Grav.).
- [16] S. W. Hawking, J. Math. Phys. **9**, 598 (1968).
- [17] S. A. Hayward, Phys. Rev. **D49**, 831 (1994).
- [18] T. Tada and S. Uehara, Phys. Rev. **D51**, 4259 (1995); Phys. Lett. B **305**, 23 (1993).
- [19] V. P. Frolov, Phys. Rev. **D46**, 5383 (1992).
- [20] J. D. Brown and J. W. York, Phys. Rev. **D47**, 1420 (1993).

- [21] J. D. Brown and J. W. York, *The path integral formulation of gravitational thermodynamics*, IFP-UNC-491, CTMP/007/NCSU, gr-qc/9405024, based on a talk presented by J. D. Brown at the conference “The Black Hole 25 Years After”, Santiago, Chile, January 1994.
- [22] T. Strobl, Phys. Rev. **D50**, 7346 (1994); P. Schaller and T. Strobl, Mod. Phys. Lett. **A9**, 3129 (1994); P. Schaller and T. Strobl, *Poisson sigma models: a generalization of gravity-Yang-Mills systems in two dimensions*, hep-th/9411163; P. Schaller and T. Strobl, *Quantization of field theories generalizing gravity-Yang-Mills systems on the cylinder*, gr-qc/9406027 or *Lecture Notes in Physics* **436**, p. 98-122, “Integrable Models and Strings,” eds. A. Alekseev, A. Hietamaeki, K. Huitu, A. Morozov, and A. Niemi; T. Strobl, *Poisson-structure induced field theory and models of 1+1 dimensional gravity*, Ph.D. dissertation, TU Wien, June 1994; T. Klösch and T. Strobl, *Classical and quantum gravity in 1+1 dimensions; part I: a unifying approach*, Technische Universität Wien preprint, August 1995 (to appear in Class. Quantum Grav.).
- [23] W. Kummer and P. Widerin, *Conserved quasilocal quantities and generally covariant theories in two dimensions*, preprint TUW-94-24, gr-qc/9502031, Vienna, December 1994 (to appear in Phys. Rev. **D**).
- [24] W. Kummer and S. R. Lau, in preparation.
- [25] M. Varadarajan, “Classical and quantum dynamics of 2d vacuum dilatonic black holes,” gr-qc/9508039, University of Utah preprint, August 1995.
- [26] A. Strominger, *Les Houches lectures on black holes*, Lectures presented at the 1994 Les Houches Summer School “Fluctuating Geometries in Statistical Mechanics and Field Theory”, gr-qc/9501071, January 1995.